

ENLARGEMENTS OF SCHEMES

LARS BRÜNJES, CHRISTIAN SERPÉ

ABSTRACT. In this article we use our constructions from [BS05] to lay down some foundations for the application of A. Robinson's nonstandard methods to modern Algebraic Geometry. The main motivation is the search for another tool to transfer results from characteristic zero to positive characteristic and vice versa. We give applications to the resolution of singularities and weak factorization.

1. INTRODUCTION

The difficulty of many problems about algebraic varieties depends on the characteristic of the base field. Resolution of singularities (proved in characteristic zero, open in characteristic p) and Grothendieck's standard conjecture on the rationality of Künneth components (proved over finite fields, open in characteristic zero) are prominent examples. This is mostly due to the fact that some tools — like transcendental methods — are only available in characteristic zero while others — like Frobenius morphisms — only exist in characteristic p .

A link between the apparently so different worlds of characteristic zero and characteristic p is provided by internal fields of *infinite* characteristic, for example the $*$ finite field $*\mathbb{Z}/P$, where $*\mathbb{Z}$ is an enlargement of \mathbb{Z} and $P \in *\mathbb{Z}$ is an infinite prime:

Let Φ be a *first order statement* in the *language of fields*. If Φ is true for all fields of characteristic zero, it is in particular true for $*\mathbb{Z}/P$ (which externally has characteristic zero), so by the permanence principle it is true for \mathbb{F}_p for infinitely many *finite* primes $p \in \mathbb{Z}$. If, on the other hand, Φ is true for almost all \mathbb{F}_p , it is also true for $*\mathbb{Z}/P$, a field of characteristic zero.

Unfortunately, being first order is a strong condition in whose absence the above reasoning fails, and the language of fields is ill adapted to dealing with schemes, sheaves and cohomology in Grothendieck's modern language of Algebraic Geometry.

Building on our paper [BS05], we therefore use the notion of *enlargement of categories* to establish a more flexible method of transferring properties from characteristic zero to characteristic p and vice versa in the framework of schemes:

Starting from a category \mathcal{B} of rings, we consider the fibred category $Sch_{\mathcal{B}}^{\text{fp}}/\mathcal{B}$ of finitely presented schemes over objects of \mathcal{B} and enlarge it to get the category of $*$ schemes $*Sch_{\mathcal{B}}^{\text{fp}}$, fibred over $*\mathcal{B}$. Here the main point is the following:

An object A of $*\mathcal{B}$ is also an ordinary ring, and we can consider the category Sch_A^{fp} of finitely presented schemes over A . The notion of scheme is *not* first order, so an object X of Sch_A^{fp} is *not* an A - $*$ scheme. Nevertheless, X is given by finitely many equations in finitely many unknowns, and these define a $*$ scheme NX over A (in fact, we construct a canonical fibred functor from Sch_A^{fp} to $*Sch_{*A}^{\text{fp}}$, which turns out to be a fibred Kan extension and is therefore unique up to unique isomorphism). Similarly, any finitely presented \mathcal{O}_X -module \mathcal{F} defines a $*$ finitely-presented \mathcal{O}_{NX} - $*$ module given by "the same" presentation. For modules, there is even a canonical functor S in the opposite direction, sending \mathcal{O}_{NX} - $*$ modules to \mathcal{O}_X -modules, and the functors N and S turn out to have many nice properties.

The main part of our paper is devoted to proving that many properties of X (like for example being smooth or proper) translate into corresponding properties of NX . — Let us stress the fact that this is *not* simply an application of the transfer principle, because the standard universe does not contain A and X and is thus not applicable.

Especially in the case where A is a *field*, properties of NX often also imply corresponding properties of X — for example, NX is $*$ irreducible respectively $*$ integral if *and only if* X is irreducible respectively integral.

Furthermore, we can give criteria (mostly of cohomological nature) for whether a given $*$ scheme or $*$ module lies in the essential image of N , thus enabling us to deduce the existence of schemes and modules

with certain properties from the existence of \ast schemes and \ast modules with the corresponding properties (note that there are many \ast scheme which do not lie in the essential image of N , for example \ast schemes of \ast finite but infinite \ast dimension and \ast schemes given by equations of \ast finite but infinite \ast degree).

At this point, let us mention Angus Macintyre's "many sorted" approach to the application of Model Theory to Algebraic Geometry in [Maci00], where he considers ultraproducts of varieties (and algebraic cycles) of fixed complexity. Though a direct comparison between Macintyre's approach and ours is difficult due to the different languages used, \ast schemes respectively \ast schemes in the essential image of N correspond to ultraproducts of varieties of arbitrary respectively bounded complexity.

The announced method of transfer between characteristic zero and characteristic p now roughly works as follows: Let Φ be a statement of schemes. Assume first that Φ holds in characteristic zero, and consider a class C of \ast schemes over \ast fields which lie in the essential image of N (i.e. a class of "bounded complexity", for example the class of \ast projective \ast schemes whose \ast dimension and \ast degree is bounded by a finite number). If k is a \ast field in $\ast\mathcal{B}$ of infinite \ast characteristic, Φ holds for schemes over k (which has characteristic zero as a field), and using properties of N , it will often be possible to show that $\ast\Phi$ then holds for \ast schemes in C , hence Φ holds for (certain) schemes over fields of *finite* characteristic (by the permanence principle). — We will give two applications of this method, namely to the problems of resolution of singularities and of weak factorization in characteristic p .

If, on the other hand, Φ holds for schemes in characteristic p , by transfer $\ast\Phi$ holds for \ast schemes over \ast fields k in $\ast\mathcal{B}$ of infinite \ast characteristic, so if X is a scheme over k , $\ast\Phi$ holds for NX . Again, using properties of N , it will often be possible to use this fact to prove that Φ holds for X , a scheme in characteristic zero. For example, if the (modified) Jacobian conjecture was proven in characteristic p , this method, combined with an easy application of the Lefschetz principle, would imply the Jacobian conjecture over \mathbb{Q} .

In subsequent papers, we plan to define similar functors N for K -theory, cycles and étale cohomology, and even though we demonstrate the usefulness of our method as it stands in the present paper (and it will not be hard to find other applications along similar lines), our main motivation for this paper is to lay the ground for that future work, from which we hope to gain new insights into the theory of algebraic cycles over varieties in characteristic zero and characteristic p .

The paper is organized as follows: In the second section we give basic definitions; in particular we define the fibration $\mathcal{Sch}_{\ast\mathcal{B}}^{\text{fp}}/\mathcal{B}$ of finitely presented schemes over a small category of rings \mathcal{B} and consider the enlargement $\ast\mathcal{Sch}_{\ast\mathcal{B}}^{\text{fp}}/\ast\mathcal{B}$.

In the third section we relate schemes and \ast schemes. For that, we define a functor $N : \mathcal{Sch}_{\ast\mathcal{B}}^{\text{fp}}/\ast\mathcal{B} \rightarrow \ast\mathcal{Sch}_{\ast\mathcal{B}}^{\text{fp}}/\ast\mathcal{B}$ which extends the canonical functor $\mathcal{Sch}_{\ast\mathcal{B}}^{\text{fp}}/\mathcal{B} \rightarrow \ast\mathcal{Sch}_{\ast\mathcal{B}}^{\text{fp}}/\ast\mathcal{B}$. In particular, for an internal ring A , we get a functor $N : \mathcal{Sch}^{\text{fp}}/A \rightarrow \ast\mathcal{Sch}^{\text{fp}}/A$.

Section 4 discusses more properties of the functor N and shows that it respects many properties of morphism between schemes.

In section 5 we define and investigate an analogous functor N for coherent modules. That is, for a scheme X of finite presentation over an internal ring, we define a functor from coherent modules on X to \ast coherent modules on NX .

Section 6 specializes to the case where the internal ring A is actually an internal field. Mainly, we apply a theorem of van den Dries and Schmidt to show — among other things — that the functor N on modules is exact and that the functor N on schemes is compatible with Quot- and Hilbert schemes.

In section 7 we show that N is compatible with higher direct images of coherent sheaves for proper morphisms (the proof of this is similar to the proof of the theorem on formal functions in Algebraic Geometry). One main application of this theorem is that N is fully faithful on coherent modules and induces an injection on Picard groups.

Section 8 shows that it is possible to define a kind of shadow map for varieties over an internal valued field with locally compact completion.

In section 9 finally we give two standard applications of the theory: First we reprove a result on resolution of singularities in characteristic p by Eklof, and second we show a similar result for the factorization of birational morphisms.

2. BASIC DEFINITIONS

Let $\mathcal{R}ings$ be the category of rings, let $\mathcal{B} \subset \mathcal{R}ings$ be a *small* (not necessarily full) subcategory, let \mathcal{R} be the small full subcategory of $\mathcal{R}ings$ containing every object of \mathcal{B} and (an isomorphic image of) every ring finitely presented over \mathbb{Z} or over an object of \mathcal{B} , and let \mathcal{S} be the small full subcategory of the category of schemes containing (an isomorphic image of) every scheme which is finitely presented over an object of \mathcal{R} .

Choose a universe \mathcal{U} such that \mathcal{S} is \mathcal{U} -small, and choose a superstructure \hat{M} containing \mathcal{U} (such that any \mathcal{U} -small category is \hat{M} -small — compare [BS05, A.3]).

Let $*$: $\hat{M} \rightarrow {}^*\hat{M}$ be an enlargement. Since \mathcal{S} is \hat{M} -small, so are \mathcal{B} and \mathcal{R} , and we can consider the enlargements ${}^*\mathcal{B} \subseteq {}^*\mathcal{R}$ and ${}^*\mathcal{S}$, all ${}^*\hat{M}$ -small categories, where ${}^*\mathcal{B}$ and ${}^*\mathcal{R}$ can be thought of as categories of (internal) rings with (internal) ring homomorphisms as morphisms (compare [BS05, 4.7]).

We call objects of ${}^*\mathcal{R}$ **rings* and objects of ${}^*\mathcal{S}$ **schemes*.

Define \mathcal{Sch} to be the category whose objects are morphisms $X \rightarrow \text{Spec}(S)$ (with X an arbitrary scheme and S an arbitrary ring) and whose morphisms $[X' \xrightarrow{\pi_{X'}} \text{Spec}(S')] \rightarrow [X \xrightarrow{\pi_X} \text{Spec}(S)]$ are pairs $\langle X' \xrightarrow{f} X, S \xrightarrow{\varphi} S' \rangle$ such that the following square commutes:

$$\begin{array}{ccc} X' & \xrightarrow{f} & X \\ \pi_{X'} \downarrow & & \downarrow \pi_X \\ \text{Spec}(S') & \xrightarrow{\text{Spec}(\varphi)} & \text{Spec}(S). \end{array}$$

(If the morphism $X \rightarrow \text{Spec}(S)$ is understood, we often denote the object $X \rightarrow \text{Spec}(S)$ by X/S or — if S is understood as well — simply by X .)

Projection onto the second component defines a functor $\mathcal{Sch} \rightarrow \mathcal{R}ings^{\text{op}}$ which is obviously a bifibration: For a ring homomorphism $\varphi : S \rightarrow S'$, inverse and direct image are given by

$$\begin{aligned} \varphi^*[X \rightarrow \text{Spec}(S)] &= [X \times_S S' \rightarrow \text{Spec}(S')] & \text{and} \\ \varphi_*[X' \rightarrow \text{Spec}(S')] &= [X' \rightarrow \text{Spec}(S') \xrightarrow{\text{Spec}(\varphi)} \text{Spec}(S)]. \end{aligned}$$

The fibre over a ring S is obviously the category \mathcal{Sch}_S of S -schemes.

Let $\mathcal{Sch}^{\text{fp}}$ be the full subcategory of \mathcal{Sch} consisting of morphisms $X \rightarrow \text{Spec}(S)$ with X a *finitely presented* S -scheme. Then $\mathcal{Sch}^{\text{fp}}$ is a subfibration of \mathcal{Sch} over $\mathcal{R}ings$ (but no longer a bifibration, because for a ring homomorphism $S \rightarrow S'$, not every finitely presented S' -scheme will in general be finitely presented as an S -scheme). Of course, the fibre over a ring S is the category $\mathcal{Sch}_S^{\text{fp}}$ of finitely presented S -schemes.

For an arbitrary subcategory \mathcal{C} of $\mathcal{R}ings$, we can form the pullbacks of $\mathcal{Sch} \rightarrow \mathcal{R}ings^{\text{op}}$ and $\mathcal{Sch}^{\text{fp}} \rightarrow \mathcal{R}ings^{\text{op}}$ along $\mathcal{C}^{\text{op}} \rightarrow \mathcal{R}ings^{\text{op}}$, and we denote the resulting bifibration respectively fibration over \mathcal{C}^{op} by $\mathcal{Sch}_{\mathcal{C}}$ respectively $\mathcal{Sch}_{\mathcal{C}}^{\text{fp}}$.[†]

Applying this to $\mathcal{C} := \mathcal{B}$ and $\mathcal{C} := {}^*\mathcal{B}$, we get bifibrations $\mathcal{Sch}_{\mathcal{B}} \rightarrow \mathcal{B}^{\text{op}}$ and $\mathcal{Sch}_{*\mathcal{B}} \rightarrow {}^*\mathcal{B}^{\text{op}}$ and fibrations $\mathcal{Sch}_{\mathcal{B}}^{\text{fp}} \rightarrow \mathcal{B}^{\text{op}}$ and $\mathcal{Sch}_{*\mathcal{B}}^{\text{fp}} \rightarrow {}^*\mathcal{B}^{\text{op}}$.

Since the fibrations $\mathcal{Sch}_{\mathcal{R}}^{\text{fp}} \rightarrow \mathcal{R}^{\text{op}}$ and $\mathcal{Sch}_{\mathcal{B}}^{\text{fp}} \rightarrow \mathcal{B}^{\text{op}}$ are obviously \hat{M} -small, we can consider their enlargements

$$\begin{array}{ccc} {}^*\mathcal{Sch}_{*\mathcal{B}}^{\text{fp}} & \hookrightarrow & {}^*\mathcal{Sch}_{*\mathcal{R}}^{\text{fp}} \\ \downarrow & & \downarrow \\ {}^*\mathcal{B}^{\text{op}} & \hookrightarrow & {}^*\mathcal{R}^{\text{op}} \end{array}$$

which are again fibrations (compare [BS05, 7.3]), whose fibres we denote by ${}^*\mathcal{Sch}_S^{\text{fp}}$ for objects S of ${}^*\mathcal{R}$.

[†]When we view \mathcal{Sch} and $\mathcal{Sch}^{\text{fp}}$ as pseudo-functors from $\mathcal{R}ings^{\text{op}}$ to the category of categories, then \mathcal{Sch}_S and $\mathcal{Sch}_S^{\text{fp}}$ are just the restrictions of these functors to \mathcal{C}^{op} .

2.1. Definition.

- (i) For a ${}^*\mathcal{R}$ -ring S , we call the category ${}^*\mathcal{A}lg_S := {}^*\mathcal{R} \setminus S$ of objects under S the category of S - ${}^*\mathcal{A}lgebras$.
- (ii) By transfer we have a functor ${}^*\mathrm{Spec} : {}^*\mathcal{R}^{\mathrm{op}} \rightarrow {}^*\mathcal{S}$ from ${}^*\mathcal{R}$ -rings to ${}^*\mathcal{S}$ -schemes, and we call ${}^*\mathcal{S}$ -schemes in the essential image of this functor ${}^*\mathcal{A}ffine$.
- (iii) For a ${}^*\mathcal{S}$ -scheme X , we call the category ${}^*\mathcal{S}ch_X := {}^*\mathcal{S} / X$ of objects over X the category of X - ${}^*\mathcal{S}chemes$ or — if $X = {}^*\mathrm{Spec}(A)$ is ${}^*\mathcal{A}ffine$ — the category ${}^*\mathcal{S}ch_A$ of A - ${}^*\mathcal{S}chemes$.
- (iv) Let \mathbf{P} be a property of rings (schemes, morphisms of rings, morphisms of schemes). When considering \mathbf{P} as a predicate on the set of objects of \mathcal{R} (of objects of \mathcal{S}, \dots), we get a predicate ${}^*\mathbf{P}$ on the set of objects of ${}^*\mathcal{R}$ (of objects of ${}^*\mathcal{S}, \dots$), i.e. a property of ${}^*\mathcal{R}$ -rings (${}^*\mathcal{S}$ -schemes, morphisms of ${}^*\mathcal{R}$ -rings, morphisms of ${}^*\mathcal{S}$ -schemes).

2.2. Remark. It follows immediately from transfer that objects of ${}^*\mathcal{S}ch_{*S}^{\mathrm{fp}}$ are morphisms of ${}^*\mathcal{S}$ -schemes $X \rightarrow {}^*\mathrm{Spec}(S)$, where S is a ${}^*\mathcal{R}$ -ring and X is a ${}^*\mathcal{S}$ -scheme. Morphisms $[X' \rightarrow {}^*\mathrm{Spec}(S')] \rightarrow [X \rightarrow {}^*\mathrm{Spec}(S)]$ are pairs $\langle f, \varphi \rangle$ with $f \in \mathrm{Mor}_{*S}(X', X)$ and $\varphi \in \mathrm{Mor}_{*\mathcal{R}}(S, S')$ such that the following square commutes in ${}^*\mathcal{S}$:

$$\begin{array}{ccc} X' & \xrightarrow{f} & X \\ \downarrow & & \downarrow \\ {}^*\mathrm{Spec}(S') & \xrightarrow{{}^*\mathrm{Spec}(\varphi)} & {}^*\mathrm{Spec}(S). \end{array}$$

In particular, for a ${}^*\mathcal{R}$ -ring S , the fibre ${}^*\mathcal{S}ch_S^{\mathrm{fp}}$ is the full subcategory of the category of S - ${}^*\mathcal{S}chemes$ defined in 2.1 consisting only of ${}^*\mathcal{F}inite$ presented S - ${}^*\mathcal{S}chemes$.

2.3. Definition. Consider the functor $\mathrm{Pol} : \mathbb{N}_0 \times \mathcal{R} \rightarrow \mathcal{R}$ (where \mathbb{N}_0 is the category associated to the partially ordered set (\mathbb{N}_0, \leq)), sending a pair (n, S) to the polynomial ring $S[X_1, \dots, X_n]$. This is a functor between \hat{M} -small categories, so we can enlarge it to a functor ${}^*\mathrm{Pol} : {}^*\mathbb{N}_0 \times {}^*\mathcal{R} \rightarrow {}^*\mathcal{R}$. For a (not necessarily finite) natural number $n \in {}^*\mathbb{N}_0$ and a ${}^*\mathcal{R}$ -ring S , we denote ${}^*\mathrm{Pol}(n, S)$ by $S^*[X_1, \dots, X_n]$ and call it the ${}^*\mathcal{P}olynomial$ ring over S in n unknowns.

2.4. Remark. Let (n, S) be an object of ${}^*\mathbb{N}_0 \times {}^*\mathcal{R}$ as above.

- (i) The morphism ${}^*\mathrm{Pol}(0 \leq n, \mathbb{1}_S) : S = S^*[] \rightarrow S^*[X_1, \dots, X_n]$ canonically turns $S^*[X_1, \dots, X_n]$ into an S - ${}^*\mathcal{A}lgebra$.
- (ii) It is easy to see that $S^*[X_1, \dots, X_n]$ has the following explicit description when viewed as an internal ring: Elements are internal ${}^*\mathcal{F}inite$ S -linear combinations of ${}^*\mathcal{M}onomials$ in n unknowns, i.e. of internal products of the form $X_1^{d_1} \cdots X_n^{d_n}$ with exponents $d_i \in {}^*\mathbb{N}_0$. These elements are added and multiplied in the obvious way.
- (iii) Transfer immediately shows that $S^*[X_1, \dots, X_n]$ has the following universal property: If T is an S - ${}^*\mathcal{A}lgebra$ and if (t_1, \dots, t_n) is an internal family of elements of T , then there is a unique morphism of S - ${}^*\mathcal{A}lgebras$ from $S^*[X_1, \dots, X_n]$ to T which sends X_i to t_i for all i .
- (iv) Let n be a ${}^*\mathcal{F}inite$ natural number. Then by the universal property of usual polynomial rings, we have a canonical morphism of S -algebras (not S - ${}^*\mathcal{A}lgebras$) $S[X_1, \dots, X_n] \rightarrow S^*[X_1, \dots, X_n]$ which sends X_i to X_i . This map is easily seen to be injective, but is (for $n \geq 1$) not bijective: For example, for an infinite $h \in {}^*\mathbb{N}_0$, the monomial X_1^h is obviously not contained in the image.

2.5. Definition. Let X be a ${}^*\mathcal{S}$ -scheme, and let n be a ${}^*\mathcal{N}atural$ number. We define the n -dimensional ${}^*\mathcal{A}ffine$ space over X as the X - ${}^*\mathcal{S}cheme$ $X^* \times_{*\mathbb{Z}} {}^*\mathbb{Z}^*[X_1, \dots, X_n]$ (note that the fibre product exists by transfer).

2.6. Remark. For every scheme X and every natural number $n \in \mathbb{N}_0$, we have the finitely presented X -scheme $\mathbb{P}_X^n = \mathbb{P}_{\mathbb{Z}}^n \times_{\mathbb{Z}} X$, the n -dimensional projective space over X , which is covered by $(n+1)$ copies of \mathbb{A}_X^n , glued together by certain universal morphisms.

By transfer, for every \ast scheme X and every \ast natural number $n \in \ast\mathbb{N}_0$, we get a \ast finitely presented X -scheme $\ast\mathbb{P}_X^n$, covered by $(n+1)$ copies of $\ast\mathbb{A}_X^n$, the n -dimensional \ast projective space over X .

If n is finite, then these \ast affine spaces are glued together by the enlargements of the corresponding morphisms from the standard world.

2.7. Definition. If S is a ring in \mathcal{R} , and if $\{f_1, \dots, f_m\} \subseteq S$ is a finite set of elements, then the category of S -algebras $A \in \text{Ob}(\mathcal{R})$ with $f_1 = \dots = f_m = 0 \in A$ has an initial object, namely the S -algebra $S/(f_1, \dots, f_m)$ (which is obviously finitely presented).

It follows by transfer that for every \ast ring S and any \ast finite internal subset $\{f_1, \dots, f_m\} \subseteq S$, there is a S - \ast algebra $S/(f_1, \dots, f_m)$ which is initial in the category of S - \ast algebras in which the f_i are mapped to zero. — We call $S/(f_1, \dots, f_m)$ the \ast factor ring of S with respect to the \ast ideal $\ast(f_1, \dots, f_m)$.[†]

2.8. Remark. Let S be a \ast ring, and let (f_1, \dots, f_m) be an ideal of S with m finite. Then it follows by easy transfer that

$$\ast(f_1, \dots, f_m) = (f_1, \dots, f_m) \cdot S \subseteq S.$$

3. RELATING SCHEMES AND \ast SCHEMES

Let A be a \ast ring in $\ast\mathcal{B}$. On the one hand, when considering A simply as a ring, we have the category Sch_A^{fp} of finitely presented A -schemes. On the other hand, we have the category $\ast\text{Sch}_A^{\text{fp}}$ of \ast finitely presented \ast schemes over A .

Intuitively, every finitely presented A -scheme determines a \ast finitely presented A - \ast scheme which is "defined by the same relations", and every morphism between finitely presented A -schemes gives a morphism between the associated A - \ast schemes.

In this section, we want to make this intuition precise by defining a morphism $N : \text{Sch}_{\ast\mathcal{B}}^{\text{fp}} \rightarrow \ast\text{Sch}_{\ast\mathcal{B}}^{\text{fp}}$ of fibrations over $\ast\mathcal{B}^{\text{op}}$. In particular, by restricting to the fibre over A , this then gives us the desired functor $\text{Sch}_A^{\text{fp}} \rightarrow \ast\text{Sch}_A^{\text{fp}}$.

3.1. Lemma. Let $\varphi : A \rightarrow B$ be a ring homomorphism in \mathcal{R} . Then the diagram

$$(1) \quad \begin{array}{ccc} A & \xrightarrow{\varphi} & B \\ \ast \downarrow & & \downarrow \ast \\ \ast A & \xrightarrow{\ast\varphi} & \ast B \end{array}$$

commutes in $\mathcal{R}\text{ings}$.

Proof. This follows immediately from elementary properties of enlargements. **q.e.d.**

3.2. Proposition/ Definition. Let A be an object of \mathcal{R} .

(i) Let $B = A[X_1, \dots, X_n]/(f_1, \dots, f_m)$ be a finitely presented A -algebra. Then

$$\ast B = \ast A[X_1, \dots, X_n]/\ast(f_1, \dots, f_m).$$

(ii) Let $\mathcal{A}\mathcal{G}_A$ respectively $\mathcal{A}\mathcal{G}_A^{\text{fp}}$ denote the category of A -algebras respectively finitely presented A -algebras. The canonical functors

$$\begin{aligned} (\mathcal{A}\mathcal{G}_A^{\text{fp}})^{\text{op}} \times \ast\mathcal{A}\mathcal{G}_{\ast A} &\longrightarrow \text{Sets} \\ (B, C) &\longmapsto \begin{cases} \text{Mor}_{\ast\mathcal{A}\mathcal{G}_{\ast A}}(\ast B, C) \\ \text{Mor}_{\mathcal{A}\mathcal{G}_A}(B, C), \end{cases} \end{aligned}$$

[†]By transfer, it is obvious that a \ast ideal of a \ast ring S is in particular an ideal of S .

induced by $*$: $\mathcal{Alg}_A^{\text{fp}} \rightarrow {}^*\mathcal{Alg}_{*A}$ and the forgetful functor ${}^*\mathcal{Alg}_{*A} \rightarrow \mathcal{Alg}_A$, are canonically isomorphic via

$$\tau_{A,B,C} : \text{Mor}_{{}^*\mathcal{Alg}_{*A}}({}^*B, C) \longrightarrow \text{Mor}_{\mathcal{Alg}_A}(B, C), \quad [{}^*B \xrightarrow{\varphi} C] \mapsto [B \xrightarrow{*} {}^*B \xrightarrow{\varphi} C].$$

Proof. By transfer, 2.4(iii) and 2.7, both *B and ${}^*A[X_1, \dots, X_n]/(f_1, \dots, f_m)$ have the same universal property in the category of *A -algebras, which proves (i).

To show (ii), we must first check that $\tau_{A,B,C}$ is indeed functorial in the arguments B and C . For argument C this is trivial, and for argument B it follows immediately from 3.1.

To see that $\tau_{A,B,C}$ is a bijection, let $B = A[X_1, \dots, X_n]/(f_1, \dots, f_m)$. Then

$$\begin{aligned} \text{Mor}_{{}^*\mathcal{Alg}_{*A}}({}^*B, C) &\stackrel{(i)}{=} \text{Mor}_{{}^*\mathcal{Alg}_{*A}}({}^*A[X_1, \dots, X_n]/(f_1, \dots, f_m), C) \\ &\stackrel{2.4(iii), 2.7}{=} \left\{ (c_1, \dots, c_n) \in C^m \mid \forall i \in \{1, \dots, m\} : f_i(c_1, \dots, c_n) = 0 \in C \right\} = \text{Mor}_{\mathcal{Alg}_A}(B, C), \end{aligned}$$

where this identification of the two sets is obviously just given by $\tau_{A,B,C}$. **q.e.d.**

3.3. Definition. For every ring A in \mathcal{R} , base change along the (external) ring homomorphism $*$: $A \rightarrow {}^*A$ defines a functor $T : \mathcal{Sch}_A \rightarrow \mathcal{Sch}_{*A}$ (which respects schemes of finite presentation), and if $\varphi : A \rightarrow A'$ is a ring homomorphism, the diagram

$$\begin{array}{ccc} \mathcal{Sch}_{A'} & \xleftarrow{\varphi^*} & \mathcal{Sch}_A \\ T \downarrow & & \downarrow T \\ \mathcal{Sch}_{*A'} & \xleftarrow{(*\varphi)^*} & \mathcal{Sch}_{*A} \end{array}$$

commutes because of 3.1. Consequently, we get "base change"-functors T of fibrations

$$\begin{array}{ccc} \mathcal{Sch}_{\mathcal{R}}^{\text{fp}} & \xrightarrow{T} & \mathcal{Sch}_{*\mathcal{R}}^{\text{fp}} \\ \downarrow & & \downarrow \\ \mathcal{Sch}_{\mathcal{R}} & \xrightarrow{T} & \mathcal{Sch}_{*\mathcal{R}} \\ \downarrow & & \downarrow \\ \mathcal{R}^{\text{op}} & \xrightarrow{*} & {}^*\mathcal{R}^{\text{op}}. \end{array}$$

For every ring A in \mathcal{R} , base change along $\text{Spec}({}^*A) \xrightarrow{\text{Spec}(*)} \text{Spec}(A)$ defines for every A -scheme X a morphism $\rho_X : TX \rightarrow X$ of schemes which is clearly functorial, i.e. the ρ_X define a 2-morphism ρ of fibrations as follows:

$$\begin{array}{ccccc} & & T & \rightarrow & \mathcal{Sch}_{*\mathcal{R}} \hookrightarrow \\ & \nearrow & & & \downarrow \rho \\ \mathcal{Sch}_{\mathcal{R}} \hookrightarrow & & & & \mathcal{Sch} \\ & \searrow & & & \downarrow \\ & & * & \rightarrow & {}^*\mathcal{R}^{\text{op}} \hookrightarrow \\ & \nearrow & & & \downarrow \rho \\ \mathcal{R}^{\text{op}} \hookrightarrow & & & & \mathcal{Rings}^{\text{op}} \end{array}$$

3.4. Theorem. There is an essentially unique functor $N : \mathcal{S}ch_{*\mathcal{R}}^{\text{fp}} \rightarrow *\mathcal{S}ch_{*\mathcal{R}}^{\text{fp}}$ of fibrations over $*\mathcal{R}^{\text{op}}$ such that the following diagram of fibrations commutes:

$$(2) \quad \begin{array}{ccccc} & & * & & \\ & \searrow & & \nearrow & \\ \mathcal{S}ch_{\mathcal{R}}^{\text{fp}} & \xrightarrow{T} & \mathcal{S}ch_{*\mathcal{R}}^{\text{fp}} & \cdots \xrightarrow{N} & *\mathcal{S}ch_{*\mathcal{R}}^{\text{fp}} \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{R}^{\text{op}} & \xrightarrow{*} & *\mathcal{R}^{\text{op}} & \xlongequal{\quad} & *\mathcal{R}^{\text{op}} \end{array}$$

In particular, by restriction to $*\mathcal{B}$, we get a canonical functor $N : \mathcal{S}ch_{*\mathcal{B}}^{\text{fp}} \rightarrow *\mathcal{S}ch_{*\mathcal{B}}^{\text{fp}}$ of fibrations over $*\mathcal{B}^{\text{op}}$.

Proof. Let A be a $*$ -ring, and let X be a scheme of finite presentation over A . According to [EGA IV₃, 8.9.1], there exist a subring $A_0 \subseteq A$, finitely generated over \mathbb{Z} , and a finitely generated (and hence finitely presented) A_0 -scheme X_0 , such that $X_0 \times_{A_0} A$ is isomorphic to X over A .

So A_0 is an object of \mathcal{R} , and X_0/A_0 is an object of $\mathcal{S}ch_{A_0}^{\text{fp}}$. According to 3.2(ii), we get the following cartesian diagram of schemes:

$$(3) \quad \begin{array}{ccccc} X & \xrightarrow{\quad} & T X_0 & \xrightarrow{\quad} & X_0 \\ \downarrow & & \downarrow & & \downarrow \\ \text{Spec}(A) & \xrightarrow{\text{Spec}(\tau_{\mathbb{Z}, A_0, A}^{-1}[A_0 \hookrightarrow A])} & \text{Spec}(*A_0) & \xrightarrow{\text{Spec}(*)} & \text{Spec}(A_0) \\ & \searrow & & \nearrow & \\ & \text{Spec}(A_0 \hookrightarrow A) & & & \end{array}$$

Therefore, in order to get a morphism of fibrations that makes (2) commute, we must define

$$N X \stackrel{(3)}{:=} \left(\tau_{\mathbb{Z}, A_0, A}^{-1}[A_0 \hookrightarrow A] \right)^* (N T X_0) \stackrel{(2)}{:=} \left(\tau_{\mathbb{Z}, A_0, A}^{-1}[A_0 \hookrightarrow A] \right)^* (*X_0).$$

Now let Y/S be another scheme of finite presentation, and let $f : X \rightarrow Y$ be an S -morphism. As before, there is a finitely generated ring $B_0 \subseteq A$ and a finitely presented B_0 -scheme Y_0 such that $Y \cong Y_0 \times_{B_0} A$.

Let I be the partially ordered set of finitely generated subrings of A containing both A_0 and B_0 , and put $X_C := X_0 \times_{A_0} C$ and $Y_C := Y_0 \times_{B_0} C$ for $C \in I$.

Then $A = \varinjlim_{C \in I} C$, $X = \varprojlim_{C \in I} X_C$ and $Y = \varprojlim_{C \in I} Y_C$, and by [EGA IV₃, 8.8.2] we have

$$(4) \quad \varinjlim_{C \in I} \text{Mor}_{\mathcal{S}ch_C}(X_C, Y_C) = \text{Mor}_{\mathcal{S}ch_A}(X, Y).$$

In particular, there exists a $C_0 \in I$ and a C_0 -morphism $f_0 : X_{C_0} \rightarrow Y_{C_0}$ such that $f = f_0 \times 1_A$. Therefore we get the following cartesian diagram of schemes

$$\begin{array}{ccccc} X & \xrightarrow{\quad} & T X_{C_0} & \xrightarrow{\quad} & X_{C_0} \\ \downarrow f & & \downarrow T f_0 & & \downarrow f_0 \\ Y & \xrightarrow{\quad} & T Y_{C_0} & \xrightarrow{\quad} & Y_{C_0} \\ \downarrow & & \downarrow & & \downarrow \\ \text{Spec}(A) & \xrightarrow{\text{Spec}(\tau_{\mathbb{Z}, C_0, A}^{-1}[C_0 \hookrightarrow A])} & \text{Spec}(*C_0) & \xrightarrow{\text{Spec}(*)} & \text{Spec}(C_0) \\ & \searrow & & \nearrow & \\ & \text{Spec}(C_0 \hookrightarrow A) & & & \end{array}$$

and we are forced to set

$$N f \stackrel{(3)}{:=} \left(\tau_{\mathbb{Z}, C_0, A}^{-1}[C_0 \hookrightarrow A] \right)^* (N T f_0) \stackrel{(2)}{:=} \left(\tau_{\mathbb{Z}, C_0, A}^{-1}[C_0 \hookrightarrow A] \right)^* (*f_0).$$

To check that this is well defined, let $C_1 \in I$ be another subring of A that admits a C_1 -morphism $f_1 : X_{C_1} \rightarrow Y_{C_1}$ with $f = f_1 \times 1_A$.

Using (4) again, we find a subring C_2 of A containing both C_0 and C_1 with $f_0 \times 1_{C_2} = f_1 \times 1_{C_2} : X_{C_2} \rightarrow Y_{C_2}$, and 3.2(ii) implies that the diagram

$$\begin{array}{ccccc}
{}^*C_0 & & \xrightarrow{\tau_{\mathbb{Z}, C_0, A}^{-1}[C_0 \hookrightarrow A]} & & A \\
& \searrow & & \nearrow & \\
& {}^*C_2 & \xrightarrow{\tau_{\mathbb{Z}, C_2, A}^{-1}[C_2 \hookrightarrow A]} & & A \\
& \nearrow & & \searrow & \\
{}^*C_1 & & \xrightarrow{\tau_{\mathbb{Z}, C_1, A}^{-1}[C_1 \hookrightarrow A]} & & A
\end{array}$$

commutes. Therefore we have

$$\begin{aligned}
\left(\tau_{\mathbb{Z}, C_0, A}^{-1}[C_0 \hookrightarrow A]\right)^*({}^*f_0) &= \left(\tau_{\mathbb{Z}, C_2, A}^{-1}[C_2 \hookrightarrow A]\right)^*({}^*f_0 \times 1_{C_2}) \\
&= \left(\tau_{\mathbb{Z}, C_2, A}^{-1}[C_2 \hookrightarrow A]\right)^*({}^*f_1 \times 1_{C_2}) = \left(\tau_{\mathbb{Z}, C_1, A}^{-1}[C_1 \hookrightarrow A]\right)^*({}^*f_1).
\end{aligned}$$

Thus Nf is well defined, and since this definition is obviously functorial, we get a functor $N : \mathcal{Sch}_A^{\text{fp}} \rightarrow {}^*\mathcal{Sch}_A^{\text{fp}}$ which furthermore is uniquely determined (up to isomorphism) by the conditions stated in the theorem.

It remains to show that this functor is compatible with inverse images and hence defines a morphism of fibrations $N : \mathcal{Sch}_A^{\text{fp}} \rightarrow {}^*\mathcal{Sch}_A^{\text{fp}}$ as claimed: If $\varphi : A \rightarrow A'$ is any morphism of * rings, we have to show that $N\varphi^*f = \varphi^*Nf$ (for $f : X \rightarrow Y$ as above). With $D_0 := \varphi(C_0) \subseteq A'$ we have

$$\varphi^*f = (f_0 \times 1_A) \times 1_{A'} = f_0 \times 1_{A'} = (f_0 \times 1_{D_0}) \times 1_{A'},$$

so

$$\begin{aligned}
N\varphi^*f &= \left(\tau_{\mathbb{Z}, D_0, A'}^{-1}[D_0 \hookrightarrow A']\right)^*({}^*(f_0 \times 1_{D_0})) = \left(\tau_{\mathbb{Z}, D_0, A'}^{-1}[D_0 \hookrightarrow A']\right)^*({}^*C_0 \rightarrow {}^*D_0)^*({}^*f_0) \\
&= \left({}^*C_0 \rightarrow {}^*D_0 \xrightarrow{\tau_{\mathbb{Z}, D_0, A'}^{-1}[D_0 \hookrightarrow A']} A'\right)^*({}^*f_0) \stackrel{3.2(\text{ii})}{=} \left(\tau_{\mathbb{Z}, C_0, A'}^{-1}[C_0 \rightarrow D_0 \xrightarrow{\varphi} A']\right)^*({}^*f_0) \\
&= \left(\tau_{\mathbb{Z}, C_0, A'}^{-1}[C_0 \hookrightarrow A \xrightarrow{\varphi} A']\right)^*({}^*f_0) \stackrel{3.2(\text{ii})}{=} \left({}^*C_0 \xrightarrow{\tau_{\mathbb{Z}, C_0, A}^{-1}[C_0 \hookrightarrow A]} A \xrightarrow{\varphi} A'\right)^*({}^*f_0) \\
&= \varphi^*\left(\tau_{\mathbb{Z}, C_0, A}^{-1}[C_0 \hookrightarrow A]\right)^*({}^*f_0) = \varphi^*Nf.
\end{aligned}$$

q.e.d.

3.5. Remark. The uniqueness of N in 3.4 can be made precise as follows: It is easy to see that N is a right Kan extension of * along T in the 2-category of fibrations (compare [MacL98, XII.4]), therefore enjoys a universal property and consequently is uniquely determined up to a canonical 2-isomorphism between morphisms of fibrations.

3.6. Example. Let A be a * ring, and let $B = A[X_1, \dots, X_n]/(f_1, \dots, f_m)$ be a finitely presented A -algebra. Let A_0 be the subring of A generated by the (finitely many) coefficients of the f_i . Then we can consider the f_i as elements of $A_0[X_1, \dots, X_n]$, and we have $B = A_0[X_1, \dots, X_n]/(f_1, \dots, f_m) \otimes_{A_0} A$. Hence

$$\begin{aligned}
(5) \quad N \operatorname{Spec}(B) &= \left(\tau_{\mathbb{Z}, A_0, A}^{-1}[A_0 \subseteq A]\right)^* \left[\operatorname{Spec}(A_0[X_1, \dots, X_n]/(f_1, \dots, f_m)) \right] \\
&\stackrel{3.2(\text{i})}{=} \left(\tau_{\mathbb{Z}, A_0, A}^{-1}[A_0 \subseteq A]\right)^* \left[{}^*\operatorname{Spec}({}^*A_0[{}^*X_1, \dots, {}^*X_n]/({}^*f_1, \dots, {}^*f_m)) \right] \\
&\stackrel{\text{transfer}}{=} {}^*\operatorname{Spec}(A[{}^*X_1, \dots, {}^*X_n]/({}^*f_1, \dots, {}^*f_m)).
\end{aligned}$$

In particular, for $n \in \mathbb{N}_0$ we get $N \mathbb{A}_A^n = {}^*\mathbb{A}_A^n$ and — taking $n = 0$ — $N \operatorname{Spec}(A) = {}^*\operatorname{Spec}(A)$.

3.7. Proposition. Let A be a * ring, let X be a finitely presented A -scheme, and let $n \in \mathbb{N}_0$ be a natural number. Then

$$N(\mathbb{P}_X^n \xrightarrow{\text{can}} X) = {}^*\mathbb{P}_{N X}^n \xrightarrow{\text{can}} N X.$$

Proof. We know from the proof of 3.4 that there exist a finitely generated subring A_0 of A and a finitely presented A_0 -scheme X_0 with $X = X_0 \times_{A_0} A$. Then

$$(\mathbb{P}_X^n \xrightarrow{\text{can}} X) = (\mathbb{P}_{X_0}^n \xrightarrow{\text{can}} X_0) \times 1_A,$$

and

$$N(\mathbb{P}_X^n \xrightarrow{\text{can}} X) = \left(\tau_{\mathbb{Z}, A_0, A}^{-1} [A_0 \hookrightarrow A] \right)^* [\mathbb{P}_{X_0}^n \xrightarrow{\text{can}} X_0] = \mathbb{P}_{NX}^n \xrightarrow{\text{can}} NX.$$

q.e.d.

4. PROPERTIES OF THE FUNCTOR N

Let A be a $*$ ring in $*\mathcal{B}$.

4.1. Proposition. The functor $N : \mathcal{Sch}_A^{\text{fp}} \rightarrow *\mathcal{Sch}_A^{\text{fp}}$

- (i) is left exact, i.e. commutes with finite limits;
- (ii) commutes with finite gluing data, i.e. if I is a finite set, if $\coprod_{i,j \in I} U_{ij} \rightrightarrows \coprod_{i \in I} U_i$ with U_{ij}, U_i finitely presented A -schemes is gluing data for an A -scheme X , then $\coprod_{i,j \in I} NU_{ij} \rightrightarrows \coprod_{i \in I} NU_i$ is gluing data for NX ;
- (iii) sends the empty scheme to the empty $*$ scheme;
- (iv) commutes with finite sums.

Proof. Let I be a *finite* category, and let $F : I \rightarrow \mathcal{Sch}_A^{\text{fp}}, i \mapsto X^i$ be an arbitrary functor. According to [EGA IV₃, 8.8.3], there exist a finitely generated subring A_0 of A and a functor $F_0 : I \rightarrow \mathcal{Sch}_{A_0}^{\text{fp}}, i \mapsto X_0^i$, such that $(\varprojlim_{i \in I} X_0^i) \times_{A_0} A = \varprojlim_{i \in I} X^i$. Since $*$ is exact by [BS05], and since inverse image functors in $*\mathcal{Sch}_{\mathcal{R}}^{\text{fp}}$ are left exact by transfer, we get $N(\varprojlim_{i \in I} X^i) = \varprojlim_{i \in I} NX^i$ by 3.4. Therefore (i) holds.

Now let I be a finite set, and let $\coprod_{i,j \in I} U_{ij} \rightrightarrows \coprod_{i \in I} U_i$ and X be as in (ii). By [EGA IV₃, 8.8.2, 8.10.5], there are a finitely generated subring A_0 of A and gluing data $\coprod_{i,j \in I} V_{ij} \rightrightarrows \coprod_{i \in I} V_i$, where the V_{ij} and V_i are finitely presented A_0 -schemes and where base change with $A_0 \hookrightarrow A$ gives back the original gluing data over A — let X_0 be the finitely presented A_0 -scheme defined gluing the V_i along the V_{ij} .

It follows from the construction of fibre products in [EGA I, 3.2.6.3] that base changes in the category of schemes respect gluing data. This implies firstly that $X_0 \times_{A_0} A = X$ and secondly (by transfer) that inverse image functors in $*\mathcal{Sch}_{\mathcal{R}}^{\text{fp}}$ commute with gluing data as well. Combining this with the exactness of $*$ (note that “commuting with gluing data” means commuting with certain finite colimits) completes the proof of (ii) using the same reasoning as for (i).

Let 0 denote the trivial ring, and let $\emptyset = \text{Spec}(0)$ be the empty (finitely presented) A -scheme. Then $\emptyset = [*Z \rightarrow A]^*\text{Spec}(0)$, so

$$N\emptyset \stackrel{3.4}{=} [*Z \rightarrow A]^*(N\text{Spec}(0)) \stackrel{3.6}{=} [*Z \rightarrow A]^*(*\text{Spec}(0)) \stackrel{\text{transfer}}{=} *\text{Spec}(0),$$

which is the empty $*$ scheme.

Finally, (iv) is just the special case of (ii) where all the U_{ij} are empty, and combining (ii) with (iii) immediately finishes the proof. **q.e.d.**

4.2. Remark. Combining 3.6 with 4.1(ii) provides us with an alternative description of the functor N , at least when we restrict our attention to *separated* A -schemes of finite presentation:

Every finitely presented A -scheme X admits a finite open affine covering $X = \bigcup_{i \in I} U_i$, and if X/A is separated, the intersections $U_{ij} := U_i \cap U_j$ are affine as well by [EGA I, 5.5.6]. So in this case, we can compute the NU_{ij} and NU_i using 3.6, and we know from 4.1(ii) that NX is obtained by glueing the NU_i along the NU_{ij} .

4.3. Corollary. Let G be a finitely presented (commutative) A -group scheme. Then NG is a $*$ finitely presented (commutative) A - $*$ group $*$ scheme, i.e. a (commutative) group object in $*\mathcal{Sch}_A^{\text{fp}}$.

Proof. The data defining a (commutative) group scheme structure on G can be expressed with diagrams involving only A , G , $G \times_A G$ and $G \times_A G \times_A G$, and these products are respected by N according to 4.1(i). **q.e.d.**

4.4. Proposition. Let $f : X \rightarrow Y$ be a morphism of finitely presented A -schemes, and let \mathbf{P} be one of the following properties of morphisms of schemes:

- isomorphism,
- monomorphism,
- immersion,
- open immersion,
- closed immersion,
- separated,
- surjective,
- radicial,
- affine,
- quasi-affine,
- finite,
- quasi-finite,
- proper,
- projective,
- quasi-projective.

If f has property \mathbf{P} , then $Nf : NX \rightarrow NY$ has property $^*\mathbf{P}$.

Proof. Let \mathbf{P} be one of the above properties. By [EGA IV₃, 8.8.2, 8.10.5], there exist a finitely generated ring $A_0 \subseteq A$ and a morphism $f_0 : X_0 \rightarrow Y_0$ of finitely presented A_0 -schemes such that $X_0 \times_{A_0} A = X$, $Y_0 \times_{A_0} A = Y$, $f_0 \times 1_A = f$ and such that f_0 has property \mathbf{P} .

Then $^*f_0 : ^*X_0 \rightarrow ^*Y_0$ has property $^*\mathbf{P}$, and since property $^*\mathbf{P}$ is stable under base change (by transfer, because \mathbf{P} is stable under base change), we see that $Nf \stackrel{3.4}{=} (\tau_{\mathbb{Z}, A_0, A}^{-1}[A_0 \subseteq A])^*(^*f_0)$ has property $^*\mathbf{P}$ as well. **q.e.d.**

4.5. Remark. Let X be a finitely presented A -scheme, and let $U \subseteq X$ be an open subscheme. According to [EGA IV₁, 1.6.2(i),(v)], U is a finitely presented A -scheme if and only if U is quasi-compact. It follows that NU is defined (and then a * open * subscheme of NX by 4.4) if and only if U is quasi-compact.

Note that the quasi-compact open subsets of X form a basis for the Zariski topology (since affine open sets are quasi-compact), so that there will be no harm in restricting our attention to quasi-compact open subschemes.

4.6. Corollary. Let $f : X \rightarrow Y$ be a morphism of finitely presented A -schemes, and let $U \subseteq Y$ be a quasi-compact open subscheme of Y . Then NU is an open * subscheme of NY , and

$$N(f|_{f^{-1}(U)}) = (Nf)|_{(Nf)^{-1}(NU)} \in \text{Mor}_{*_{\text{Sch}_A^{\text{fp}}}}(N(f^{-1}(U)), NU)$$

Proof. This follows immediately from the fact that N is left exact by 4.1(i) and respects open immersions by 4.4, applied to the cartesian diagram

$$\begin{array}{ccc} f^{-1}(U) & \xrightarrow{f|_{f^{-1}(U)}} & U \\ \downarrow & & \downarrow \\ X & \xrightarrow{f} & Y. \end{array}$$

q.e.d.

4.7. Corollary. Let $X = \bigcup_{i \in I} U_i$ be a *finite* (affine) covering by quasi-compact open subschemes. Then $NX = \bigcup_{i \in I} NU_i$ is a \ast open (\ast affine) \ast covering in $\ast\text{Sch}_A^{\text{fp}}$.

Proof. If the U_i are affine, the NU_i are \ast affine by example 3.6. The NU_i are open subschemes of NX by 4.4, and since

$$\coprod_{i \in I} NU_i \xrightarrow{4.1} N\left(\coprod_{i \in I} U_i\right) \xrightarrow{4.4} NX,$$

they cover NX . **q.e.d.**

4.8. Lemma. Let X be a finitely presented A -scheme, let $Y \subseteq X$ be a closed, finitely presented subscheme, and assume that the open complement $U := X \setminus Y$ is quasi-compact. Then NU is $[NX] \setminus [NY]$, the \ast complement of NY in NX .

Proof. Since the diagram

$$\begin{array}{ccc} \emptyset & \xrightarrow{\subset} & U \\ \downarrow & \square & \downarrow \\ Y & \xrightarrow{\subset} & X \end{array}$$

is cartesian, 4.1(i), (iii) imply that

$$\begin{array}{ccc} \emptyset & \xrightarrow{\subset} & NU \\ \downarrow & \square & \downarrow \\ NY & \xrightarrow{\subset} & NX \end{array}$$

is also cartesian, i.e. NU lies in $[NX] \setminus [NY]$. For the other inclusion, note that the surjectivity of $Y \coprod U \rightarrow X$ implies the \ast surjectivity of $[NY] \coprod [NU] \rightarrow NX$ by 4.1(iv) and 4.4. **q.e.d.**

Let $\varphi : R \rightarrow S$ be a ring homomorphism, let X be an R -scheme, and let Y be an S -scheme. Then it is common practice to simply write $X(Y)$ for the set of those morphisms $f : Y \rightarrow X$ of schemes that make the diagram

$$\begin{array}{ccc} Y & \xrightarrow{f} & X \\ \downarrow & & \downarrow \\ \text{Spec}(S) & \xrightarrow{\text{Spec}(\varphi)} & \text{Spec}(R) \end{array}$$

commute, thus dropping R, S and φ from the notation. In other words, when R, S and φ are understood, $X(Y)$ denotes the subset of those morphisms in Sch which project to φ in the bifibration $\text{Sch} \rightarrow \mathcal{R}\text{ings}^{\text{op}}$.

In analogy to this practice, we make the following definition:

4.9. Definition. Let X be a \ast scheme in $\ast\text{Sch}_A^{\text{fp}}$, let $\varphi : A \rightarrow B$ be a morphism of \ast rings, and let Y be a \ast scheme in $\ast\text{Sch}_B^{\text{fp}}$.

Then we denote the set of those morphisms in $\text{Mor}_{\ast\text{Sch}_A^{\text{fp}}}(Y/B, X/A)$ which are projected to φ under $\ast\text{Sch}_A^{\text{fp}} \rightarrow \ast\mathcal{R}^{\text{op}}$ by $X(Y)$ and call it the set of Y -valued points of X (where we assume that A, B and φ are understood).

In the special case $Y = \ast\text{Spec}(B)$, we put $X(B) := X(Y)$ and call $X(B)$ the set of B -valued points of X .

4.10. Remark. Let X be a finitely presented A -scheme, let $\varphi : A \rightarrow B$ be a morphism of \ast rings, and let T be a finitely presented B -scheme.

Then the functor N induces a canonical map

$$\begin{array}{ccc} X(T) & \xrightarrow{\quad N \quad} & (NX)(NT) \\ \downarrow & & \downarrow \\ \text{Mor}_{\text{Sch}_B^{\text{fp}}}(T/B, X/A) & \xrightarrow{\quad N \quad} & \text{Mor}_{\text{Sch}_B^{\text{fp}}}(NT/B, NX/A) \end{array}$$

(note that N , restricted to $X(T)$, factorizes over $(NX)(NT)$, because N is a morphism of fibrations and hence in particular a morphism of categories over ${}^*\mathcal{R}^{\text{op}}$).

Since $N\text{Spec}(B) = {}^*\text{Spec}(B)$ by 3.6, we in particular get a map $N : X(B) \rightarrow (NX)(B)$ from B -valued points of X to B valued points of NX .

4.11. Definition. As we have seen in 3.6, the functor $N : \text{Sch}_A^{\text{fp}} \rightarrow {}^*\text{Sch}_A^{\text{fp}}$ sends affine schemes to * affine schemes and thus induces a functor $\mathcal{A}lg_A^{\text{fp}} \rightarrow {}^*\mathcal{A}lg_A^{\text{fp}}$ — which we want to denote by N as well — satisfying

$$(6) \quad \forall B \in \text{Ob}(\mathcal{A}lg_A^{\text{fp}}) : N\text{Spec}(B) = {}^*\text{Spec}(NB).$$

If $B = A[X_i]/(f_j)$, then we have calculated in 3.6 that $NB = A^*[X_i]/^*(f_j)$. It follows from 2.4(iii) and 2.7 that sending X_i to X_i defines a canonical morphism of A -algebras $\sigma_B : B \rightarrow NB$, which is obviously functorial: If $\varphi : B \rightarrow C$ is a morphism of A -algebras, then

$$(7) \quad \begin{array}{ccc} B & \xrightarrow{\sigma_B} & NB \\ \varphi \downarrow & & \downarrow N\varphi \\ C & \xrightarrow{\sigma_C} & NC \end{array}$$

commutes in the category of A -algebras.

4.12. Lemma. Let k be an A - * algebra, and let B be a finitely presented A -algebra. Then the canonical map

$$(\sigma_B)_* : \text{Mor}_{{}^*\mathcal{A}lg_A}(NB, k) \longrightarrow \text{Mor}_{\mathcal{A}lg_A}(B, k), \quad [NB \xrightarrow{\varphi} k] \mapsto [B \xrightarrow{\sigma_B} NB \xrightarrow{\varphi} k]$$

is bijective.

Proof. Let $B = A[X_i]/(f_j)$. We can argue as in the proof of 3.2(ii): A morphism $\varphi : NB \rightarrow k$ in ${}^*\mathcal{A}lg_A$ is precisely given by a tuple $(x_1, \dots, x_n) \in k^n$ satisfying $f_j(x_1, \dots, x_n) = 0 \in k$ for all j , and the exact same data defines a morphism $\varphi' : B \rightarrow k$ of A -algebras. — It is clear that this identification between the two sets of morphisms is just the one given in the lemma. **q.e.d.**

4.13. Theorem. Let k be a * artinian A - * algebra, and let X be a finitely presented A -scheme. Then the canonical map $N : X(k) \rightarrow (NX)(k)$ is bijective.

Proof. We choose a finite affine open covering $X = \bigcup_{i \in I} U_i$, so that $NX = \bigcup_{i \in I} NU_i$ is a * open * affine * covering of NX by 4.7.

To prove surjectivity, let $f : {}^*\text{Spec}(k) \rightarrow X$ be an arbitrary k -valued point of X . By transfer, since k is * artinian, f factorizes over one of the NU_i , so without loss of generality, we can assume that $X = \text{Spec}(B)$ is affine.

Then $NX \stackrel{(6)}{=} {}^*\text{Spec}(NB)$, and f corresponds to a morphism $\varphi : NB \rightarrow k$ of A - * algebras which induces a morphism $\varphi' := \varphi\sigma_B : B \rightarrow k$ of A -algebras as in 4.12, hence a k -valued point $f' := \text{Spec}(\varphi')$ of X . It is clear that $Nf' = f$, so N is indeed surjective.

For injectivity, let $f, g \in X(k)$ be two k -valued points of X with $Nf = Ng \in (NX)(k)$.

If X_k denotes the inverse image of X under $A \rightarrow k$, then the canonical map $X_k(k) \rightarrow X(k)$ is a bijection, so that we can assume $A = k$ without loss of generality. As above, it follows that f factorizes

over one of the U_i , say over U_{i_0} — then Nf factorizes over NU_{i_0} . Let us assume that g does not factorize over U_{i_0} . This would imply that the following diagram of finitely presented k -schemes is cartesian:

$$\begin{array}{ccc} \emptyset & \xrightarrow{\quad} & U_{i_0} \\ \downarrow & & \downarrow \\ \mathrm{Spec}(k) & \xrightarrow{g} & X. \end{array}$$

Then 4.1(i) and (iii) imply that

$$\begin{array}{ccc} {}^*\emptyset & \xrightarrow{\quad} & NU_{i_0} \\ \downarrow & & \downarrow \\ {}^*\mathrm{Spec}(k) & \xrightarrow{Ng=Nf} & NX \end{array}$$

is cartesian as well, a contradiction to the fact that Nf factorizes over NU_{i_0} .

Therefore both f and g factorize over U_{i_0} , and we can again assume that $X = \mathrm{Spec}(B)$ is affine. But then f and g correspond to k -algebra morphisms $\varphi, \psi : B \rightarrow k$, and $Nf = Ng$ means that the induced morphisms of k - $*$ -algebras $\varphi', \psi' : NB \rightarrow k$ are the same. But then φ and ψ must be the same as well according to 4.12. **q.e.d.**

4.14. Proposition. Let $f : X \rightarrow Y$ be a morphism of finitely presented A -schemes. If f is étale (unramified, smooth), then $Nf : NX \rightarrow NY$ is $*$ étale ($*$ unramified, $*$ smooth).

Proof. First consider the case where $f : X \rightarrow Y$ is unramified. By [EGA IV₄, 17.4.2], a morphism $f : X \rightarrow Y$ of (locally) finite presentation is unramified if and only if the diagonal $\Delta_{X/Y} : X \xrightarrow{(f,f)} X \times_Y X$ is an open immersion. So in our case, $\Delta_{X/Y}$ is an open immersion, and 4.1(i) and 4.4 show that the $*$ diagonal $\Delta_{NX/NY} : NX \xrightarrow{(Nf, Nf)} NX \times_{NY} NX$ is a $*$ open immersion, hence transferring [EGA IV₄, 17.4.2] proves that Nf is $*$ unramified (since it is $*$ finitely presented by construction).

Now let $f : X \rightarrow Y$ be étale. By [EGA IV₄, 17.1.6], 4.6 and 4.7, we can assume without loss of generality that X and Y are affine and that f is given by a morphism $\varphi : B \rightarrow C$ of finitely presented A -algebras. Furthermore, by [Mil80, I.3.16], we can assume that $C = B[T_1, \dots, T_n]/(P_1, \dots, P_n)$ with $d := \det(\partial P_i / \partial T_j) \in C^\times$ and that φ is the canonical morphism, and we have to show that $N\varphi : NB \rightarrow NC = (NB)^*[T_i]/^*(P_j)$ is $*$ étale. By transfer of [Mil80, I.3.16], for this it suffices to show that $d' := * \det(*\partial P_i / *\partial T_j)$ is a $*$ unit in NC .

Since partial derivatives of polynomials and determinants of matrices are given by universal polynomials in the coefficients, it follows easily that the diagrams

$$\begin{array}{ccc} B[T_1, \dots, T_n] & \xrightarrow{\partial/\partial T_j} & B[T_1, \dots, T_n] \\ \sigma_{B[T_i]} \downarrow & & \downarrow \sigma_{B[T_i]} \\ B^*[T_1, \dots, T_n] & \xrightarrow{*\partial/*\partial T_j} & B^*[T_1, \dots, T_n] \end{array} \quad \begin{array}{ccc} B[T_1, \dots, T_n]^{n \times n} & \xrightarrow{\det} & B[T_1, \dots, T_n] \\ \sigma_{B[T_i]}^{n \times n} \downarrow & & \downarrow \sigma_{B[T_i]} \\ B^*[T_1, \dots, T_n]^{n \times n} & \xrightarrow{*\det} & B^*[T_1, \dots, T_n] \end{array}$$

commute, which implies $d' = \sigma_C(d) \in NC$. Since σ_C is a ring homomorphism, it maps units to units, so d' is a unit in NC . But being a unit is obviously a first order property, so units and $*$ units are the same thing, and we are done in the case where f is étale.

Finally, let $f : X \rightarrow Y$ be smooth. By [Mil80, 3.24], this is equivalent to the existence of a (finite) open affine covering U_i of X , such that for every i the restriction $f|_{U_i}$ factorizes as

$$\begin{array}{ccc} U_i & \xrightarrow{f|_{U_i}} & Y \\ g_i \downarrow & & \downarrow \\ \mathbb{A}_{V_i}^n & \xrightarrow{\text{can}} & V_i \end{array}$$

with g_i étale and $n \in \mathbb{N}_0$. Since the functor N respects open affine coverings by 4.7, restrictions by 4.6, open immersions by 4.4, affine spaces (over affine bases) by 3.6 and étale morphisms by the second part of the proof, transfer of [Mil80, 3.24] shows that Nf is indeed $*$ smooth. **q.e.d.**

4.15. Lemma. Let B be a finitely presented A -algebra, and let $C = B[Y_1, \dots, Y_k]/\mathcal{I}$ be a finitely presented B -algebra. Then $NC = (NB)^*[Y_j]/*\mathcal{I}$.

Proof. Let $B = A[X_1, \dots, X_n]/I$ be a finite presentation of B as an A -algebra. Then

$$NC = N\left(A[X_i, Y_j]/(I + \mathcal{I})\right) \stackrel{3.6}{=} A^*[X_i, Y_j]/*(I + \mathcal{I}) \stackrel{\text{transfer}}{=} \left(A^*[X_i]/*I\right)^*[Y_j]/*\mathcal{I} \stackrel{3.6}{=} (NB)[Y_j]/*\mathcal{I}.$$

q.e.d.

4.16. Proposition. Let B be a finitely presented A -algebra, and let C be a *finite* B -algebra. Then the canonical ring homomorphism $C \otimes_B NB \rightarrow NC$ induced by (7) is an isomorphism.

Proof. First consider the case where $C = B/I$ is a quotient of B . Then

$$C \otimes_B NB = (NB)/I \cdot NB = (NB)/^*I \stackrel{4.15}{=} NC.$$

Next let $C = B[c]/(c^n + b_{n-1}c^{n-1} + \dots + b_0)$ with $n \in \mathbb{N}_+$ and $b_0, \dots, b_{n-1} \in B$. Consider the following true statement in \hat{M} :

For every object R of \mathcal{R} and for every tuple $(r_0, \dots, r_{n-1}) \in R^n$, sending e_i to \bar{X}^{i-1} defines an isomorphism of R -modules $R^n \xrightarrow{\sim} R[X]/(X^n + r_{n-1}X^{n-1} + \dots + r_0)$.

By transfer and the fact that an isomorphism of $*$ -modules is in particular an isomorphism of modules, we get:

For every $*$ -ring R and for every tuple $(r_0, \dots, r_{n-1}) \in R^n$, sending e_i to \bar{X}^{i-1} defines an isomorphism of R -modules $R^n \xrightarrow{\sim} R^*[X]/*(X^n + r_{n-1}X^{n-1} + \dots + r_0)$.

By 4.15, we have $NC = (NB)^*[c]/*(c^n + b_{n-1}c^{n-1} + \dots + b_0)$, so we get the following commutative diagram of NB -modules:

$$\begin{array}{ccccccc} \bar{c}^{i-1} \otimes 1 & \in & C \otimes_B NB & \xrightarrow{\dots\dots\dots} & NC & \ni & \bar{c}^{i-1} \\ \uparrow & & \uparrow \wr & & \uparrow \wr & & \uparrow \\ e_i & \in & NB^n & \xrightarrow{\sim} & NB^n & \ni & e_i, \end{array}$$

and we are done in this case as well.

Now let $C = B[c]/I$. Then the element \bar{c} of C is integral over B , because C/B is finite, so there is a relation $\bar{c}^n + b_{n-1}\bar{c}^{n-1} + \dots + b_0 = 0$ in C , which means that $B \rightarrow C$ factorizes as

$$B \rightarrow \underbrace{B[c]/(c^n + b_{n-1}c^{n-1} + \dots + b_0)}_{=: C'} \xrightarrow{c \mapsto \bar{c}} C,$$

and we get

$$NC \stackrel{1. \text{ case}}{=} C \otimes_{C'} NC' \stackrel{2. \text{ case}}{=} C \otimes_{C'} C' \otimes_B NB = C \otimes_B NB.$$

Finally, in the general case, let $C = B[X_1, \dots, X_n]/I$ for an $n \in \mathbb{N}_+$. We prove the proposition by induction on n : The case $n = 1$ has been proven above, so let $C = B[X_1, \dots, X_{n+1}]/I$ for $n \geq 1$. Let C' be the subring of C generated by $\bar{X}_1, \dots, \bar{X}_n$ as a B -algebra. Then $C = C'[X_{n+1}]/\mathcal{I}$, and

$$NC \stackrel{3. \text{ case}}{=} C \otimes_{C'} NC' \stackrel{\text{induction}}{=} C \otimes_{C'} C' \otimes_B NB = C \otimes_B NB.$$

q.e.d.

5. *MODULES OVER *SCHEMES

Let $\mathcal{M}od$ be the category whose objects are pairs $\langle \mathcal{F}, X/A \rangle$, consisting of an A -scheme X and an \mathcal{O}_X -module \mathcal{G} , and whose morphisms from $\langle \mathcal{F}, X/A \rangle$ to $\langle \mathcal{G}, Y/B \rangle$ are pairs $\langle \alpha, \langle f, \varphi \rangle \rangle$ with $\langle f, \varphi \rangle$ a morphism from X/A to Y/B in $\mathcal{S}ch$ and $\alpha : f^* \mathcal{G} \rightarrow \mathcal{F}$ a morphism of \mathcal{O}_X -modules.

Projection onto the second component defines an abelian bifibration $\mathcal{M}od \rightarrow \mathcal{S}ch$ (or $\mathcal{M}od \rightarrow \mathcal{R}ings^{\text{op}}$ after composing with $\mathcal{S}ch \rightarrow \mathcal{R}ings^{\text{op}}$): For a morphism $\langle f, \varphi \rangle : X/A \rightarrow Y/B$, direct and inverse image functor are given by $\langle f, \varphi \rangle_* \langle \mathcal{F}, X/A \rangle = \langle f_* \mathcal{F}, Y/B \rangle$ respectively $\langle f, \varphi \rangle^* \langle \mathcal{G}, Y/B \rangle = \langle f^* \mathcal{G}, X/A \rangle$, and the fibre over an object X/A is the *opposite* of the category $\mathcal{M}od_X$ of \mathcal{O}_X -modules.

Let $\mathcal{M}od_{\mathcal{R}}^{\mathcal{U}} \rightarrow \mathcal{S}ch_{\mathcal{R}}^{\text{fp}}$ be the full subcategory of the pullback of this fibration along $\mathcal{S}ch_{\mathcal{R}}^{\text{fp}} \hookrightarrow \mathcal{S}ch$ consisting of \mathcal{U} -sheaves, i.e. sheaves in our chosen universe \mathcal{U} . — this is an abelian, \hat{M} -small bifibration where the opposite of each fibre has enough injective objects.

For a scheme X , denote the category of quasi-projective (respectively finitely presented) \mathcal{O}_X -modules by $QCoh_X$ (respectively $\mathcal{M}od_X^{\text{fp}}$). Recall from [EGA I, 5.2.5] that an \mathcal{O}_X -module \mathcal{F} is called *finitely presented* if for every $x \in X$, there is an open neighborhood $U \subseteq X$ of x and an exact sequence $\mathcal{O}_U^m \rightarrow \mathcal{O}_U^n \rightarrow \mathcal{F}|_U \rightarrow 0$ of \mathcal{O}_U -modules with natural numbers m and n . If X is locally noetherian, this is equivalent to \mathcal{F} being a coherent \mathcal{O}_X -module.

Let $QCoh$ (respectively $\mathcal{M}od^{\text{fp}}$) be the full subcategory of $\mathcal{M}od$ whose fibre over X/A is the opposite of $QCoh_X$ (respectively of $\mathcal{M}od_X^{\text{fp}}$).

Pulling back along $\mathcal{S}ch_{\mathcal{R}}^{\text{fp}} \rightarrow \mathcal{S}ch$ and restricting to \mathcal{U} -sheaves, we get \hat{M} -small fibrations $QCoh_{\mathcal{R}}^{\mathcal{U}}$ and $\mathcal{M}od_{\mathcal{R}}^{\text{fp}}$ over $\mathcal{S}ch_{\mathcal{R}}^{\text{fp}}$ (note that any finitely presented \mathcal{O}_X -module for X in \mathcal{S} is automatically a \mathcal{U} -sheaf). We sum up the situation in the following diagram of additive fibrations:

$$\begin{array}{ccccccc}
 \mathcal{M}od_{\mathcal{R}}^{\text{fp}} & \hookrightarrow & QCoh_{\mathcal{R}}^{\mathcal{U}} & \hookrightarrow & \mathcal{M}od_{\mathcal{R}}^{\mathcal{U}} & \hookrightarrow & \mathcal{M}od \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \mathcal{S}ch_{\mathcal{R}}^{\text{fp}} & \xlongequal{\quad} & \mathcal{S}ch_{\mathcal{R}}^{\text{fp}} & \xlongequal{\quad} & \mathcal{S}ch_{\mathcal{R}}^{\text{fp}} & \hookrightarrow & \mathcal{S}ch \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \mathcal{R}^{\text{op}} & \xlongequal{\quad} & \mathcal{R}^{\text{op}} & \xlongequal{\quad} & \mathcal{R}^{\text{op}} & \xlongequal{\quad} & \mathcal{R}ings^{\text{op}}
 \end{array}$$

The first three columns in this diagram are \hat{M} -small, and we enlarge them to get an additive fibration $*\mathcal{M}od_{\mathcal{R}}^{\text{fp}}/*\mathcal{R}$, an abelian fibration $*QCoh_{\mathcal{R}}/*\mathcal{R}$ and an abelian bifibration $*\mathcal{M}od_{\mathcal{R}}/*\mathcal{R}$.

For a $*$ -scheme X , we denote the *opposite* of the fibre of $*\mathcal{M}od_{\mathcal{R}}$ (respectively $*\mathcal{M}od_{\mathcal{R}}^{\text{fp}}$, respectively $*QCoh_{\mathcal{R}}$) over X by $*\mathcal{M}od_X$ (respectively $*\mathcal{M}od_X^{\text{fp}}$, respectively $*QCoh_X$), and we call the objects of this fibre \mathcal{O}_X - $*$ modules (respectively $*finitely\ presented\ \mathcal{O}_X$ - $*$ modules, respectively $*quasi-coherent\ \mathcal{O}_X$ - $*$ modules).

If X is $*$ locally noetherian, we also say $*coherent$ instead of $*finitely\ presented$, and $*Coh_X := *\mathcal{M}od_X^{\text{fp}}$ is an abelian category.

5.1. Lemma/ Definition. Sending $\langle \mathcal{F}, X/A \rangle$ to $\langle \rho_X^* \mathcal{F}, T X/*A \rangle$ induces a canonical morphism of additive fibrations $T : \mathcal{M}od_{\mathcal{R}}^{\text{fp}} \rightarrow \mathcal{M}od_{*\mathcal{R}}^{\text{fp}}$:

$$\begin{array}{ccc}
 \mathcal{M}od_{\mathcal{R}}^{\text{fp}} & \xrightarrow{\quad T \quad} & \mathcal{M}od_{*\mathcal{R}}^{\text{fp}} \\
 \downarrow & & \downarrow \\
 \mathcal{S}ch_{\mathcal{R}}^{\text{fp}} & \xrightarrow{\quad T \quad} & \mathcal{S}ch_{*\mathcal{R}}^{\text{fp}} \\
 \downarrow & & \downarrow \\
 \mathcal{R}^{\text{op}} & \xrightarrow{\quad * \quad} & *\mathcal{R}^{\text{op}}
 \end{array}$$

Proof. This is obvious. **q.e.d.**

5.2. Theorem. There is an (essentially) unique morphism $N : \mathcal{M}od_{*\mathcal{R}}^{\text{fp}} \rightarrow * \mathcal{M}od_{\mathcal{R}}^{\text{fp}}$ of additive fibrations over $*\mathcal{S}ch_{\mathcal{R}}$ that makes the following diagram commute:

$$\begin{array}{ccccc}
 & & * & & \\
 & \curvearrowright & & \curvearrowleft & \\
 \mathcal{M}od_{\mathcal{R}}^{\text{fp}} & \xrightarrow{T} & \mathcal{M}od_{*\mathcal{R}}^{\text{fp}} & \dashrightarrow^N & * \mathcal{M}od_{\mathcal{R}}^{\text{fp}} \\
 \downarrow & & \downarrow & & \downarrow \\
 \mathcal{S}ch_{\mathcal{R}}^{\text{fp}} & \xrightarrow{T} & \mathcal{S}ch_{*\mathcal{R}}^{\text{fp}} & \xrightarrow{N} & * \mathcal{S}ch_{\mathcal{R}}^{\text{fp}} \\
 \downarrow & & \downarrow & & \downarrow \\
 \mathcal{R}^{\text{op}} & \xrightarrow{*} & *\mathcal{R}^{\text{op}} & \xlongequal{\quad} & *\mathcal{R}^{\text{op}}
 \end{array}$$

In particular, for every $*$ ring A and every finitely presented A -scheme X , we get a canonical additive functor $N : \mathcal{M}od_X^{\text{fp}} \rightarrow * \mathcal{M}od_{NX}^{\text{fp}}$.

Proof. This follows from [EGA IV₃, 8.5.2] in the same way as 3.4 follows from [EGA IV₃, 8.8.2]. **q.e.d.**

From now on for the rest of this section, let A be a $*$ ring, and let X be a finitely presented A -scheme.

5.3. Proposition. Let $\mathcal{F} \rightarrow \mathcal{G} \rightarrow \mathcal{H} \rightarrow 0$ be a sequence in $\mathcal{M}od_X^{\text{fp}}$ which is exact in $\mathcal{M}od_X$. Then the sequence $N\mathcal{F} \rightarrow N\mathcal{G} \rightarrow N\mathcal{H} \rightarrow 0$ of $*$ finitely-presented \mathcal{O}_{NX} - $*$ modules is exact in $*\mathcal{M}od_{NX}$.

In particular, if A is noetherian (for example a $*$ field), then the functor N from coherent \mathcal{O}_X -modules to $*$ coherent \mathcal{O}_{NX} - $*$ modules is right exact.

Proof. This follows from [EGA IV₃, 8.5.6] and the construction of N . **q.e.d.**

5.4. Proposition. For $n \in \mathbb{N}_0$, we have $N \mathcal{O}_X^n = \mathcal{O}_{NX}^n$.

Proof. Since N is additive, we only have to consider the case $n = 1$. Because A is a $*$ ring, we have a canonical morphism of $*$ rings $*\mathbb{Z} \rightarrow A$ and hence a canonical morphism $f : X/A \rightarrow \text{Spec}(*\mathbb{Z})/*\mathbb{Z}$ in $\mathcal{S}ch_{*\mathcal{R}}^{\text{fp}}$. Then $\mathcal{O}_X = f^* \mathcal{O}_{\text{Spec}(*\mathbb{Z})}$, so

$$N \mathcal{O}_X \stackrel{5.2}{=} (Nf)^* N \mathcal{O}_{\text{Spec}(*\mathbb{Z})} = (Nf^*) N T \mathcal{O}_{\text{Spec}(\mathbb{Z})} = (Nf)^* \mathcal{O}_{*\text{Spec}(\mathbb{Z})} = \mathcal{O}_{NX}.$$

q.e.d.

5.5. Corollary. Let \mathcal{E} be a vector bundle of rank $n \in \mathbb{N}_0$ on X . Then $N\mathcal{E}$ is a $*$ vector bundle of rank n on NX .

Proof. This follows immediately from 5.4. **q.e.d.**

5.6. Lemma/ Definition. For an \mathcal{O}_{NX} - $*$ module \mathcal{F} , sending a quasi-compact open subscheme U of X to $\mathcal{F}(NU)$ defines an abelian sheaf $N_*\mathcal{F}$ on X . In this way, we get an additive functor N_* from $*\mathcal{M}od_{NX}$ to the category of abelian sheaves on X .

Proof. First of all, $N_*\mathcal{F}$ is clearly an abelian presheaf on the category of quasi-compact open subsets of X , because N is a functor from that category to the category of $*$ open $*$ subschemes of NX . By 4.5, such a presheaf defines a sheaf on X , provided the sheaf-condition with respect to finite, quasi-compact, open coverings is satisfied.

So let $U \subseteq X$ be quasi-compact and open, and let $U = U_1 \cup \dots \cup U_n$ be a finite, quasi-compact, open covering of U . Then by 4.7, $[NU] = [NU_1] \cup \dots \cup [NU_n]$ is a $*$ open covering of NU , which is *internal* because it is finite. By transfer, since \mathcal{F} is a \mathcal{O}_{NX} - $*$ module, we get the following exact sequence (of abelian $*$ groups):

$$0 \longrightarrow \mathcal{F}(NU) \longrightarrow \prod_{i=1}^n \mathcal{F}(NU_i) \longrightarrow \prod_{i,j=1}^n \mathcal{F}(NU_i \cap NU_j).$$

But n is finite, and finite $*$ products are simply products, so we get the following sequence of abelian groups

$$0 \longrightarrow [N_*\mathcal{F}](U) \longrightarrow \prod_{i=1}^n [N_*\mathcal{F}](U_i) \longrightarrow \prod_{i,j=1}^n [N_*\mathcal{F}](U_i \cap U_j),$$

which is just the sheaf condition we wanted to prove, so $N_*\mathcal{F}$ is indeed an abelian sheaf on X .

Finally, since N is a functor, we really get an additive functor N_* as desired. **q.e.d.**

5.7. Definition. Since $N_*\mathcal{O}_{NX}$ is a sheaf of rings on X by 5.6, we get a ringed space

$$\hat{X} := (X, \mathcal{O}_{\hat{X}}) := (X, N_*\mathcal{O}_{NX}),$$

and from now on, we want to consider N_* as an additive functor from ${}^*\mathcal{M}od_{NX}$ to $\mathcal{M}od_{\hat{X}}$.

If $U = \text{Spec}(B)$ is an affine, open subscheme of X , then we have a canonical morphism of A -algebras

$$\mathcal{O}_X(U) = B \xrightarrow{\sigma_B} NB = \mathcal{O}_{NX}(NU) = \mathcal{O}_{\hat{X}}(U),$$

which is functorial in U by (7), i.e. we get a morphism of sheaves of rings $\sigma : \mathcal{O}_X \rightarrow \mathcal{O}_{\hat{X}}$ on X and hence a canonical morphism of ringed spaces $\sigma^* : \hat{X} \rightarrow X$, which in turn defines a canonical additive functor $\sigma_* : \mathcal{M}od_{\hat{X}} \rightarrow \mathcal{M}od_X$. We denote the composition

$${}^*\mathcal{M}od_{NX} \xrightarrow{N_*} \mathcal{M}od_{\hat{X}} \xrightarrow{\sigma_*} \mathcal{M}od_X$$

by S .

5.8. Proposition. The functors $N_* : {}^*\mathcal{M}od_{NX} \rightarrow \mathcal{M}od_{\hat{X}}$ and $S : {}^*\mathcal{M}od_{NX} \rightarrow \mathcal{M}od_X$ are left exact, and their restrictions to ${}^*Q\mathcal{C}oh_{NX}$ are exact and faithful.

Proof. The functor $\sigma_* : \mathcal{M}od_{\hat{X}} \rightarrow \mathcal{M}od_X$ is exact and faithful, because it is the identity functor on the underlying abelian sheaves, so if S is left exact respectively exact and faithful, so is N_* .

Let $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ be an exact sequence of \mathcal{O}_{NX} - $*$ modules. If U is a $*$ open $*$ subscheme of NX , then by transfer the sequence

$$0 \longrightarrow \mathcal{F}'(U) \longrightarrow \mathcal{F}(U) \longrightarrow \mathcal{F}''(U)$$

is exact (in the category of internal $\mathcal{O}_{NX}(U)$ -modules and hence in particular in the category of abelian groups), which proves that S is left exact.

Now let \mathcal{F}' , \mathcal{F} and \mathcal{F}'' be $*$ quasi-coherent. Let $x \in X$ be an arbitrary point, and let $t_x \in [S\mathcal{F}'']_x$ be an arbitrary element in the stalk. There is an affine open subscheme U of X with a local section $t_U \in [S\mathcal{F}''](U)$ which represents t_x . Since \mathcal{F}' , \mathcal{F} and \mathcal{F}'' are $*$ quasi-coherent and since NU is $*$ affine, it follows by transfer that

$$0 \longrightarrow \mathcal{F}'(NU) \longrightarrow \mathcal{F}(NU) \longrightarrow \mathcal{F}''(NU) \longrightarrow 0$$

is exact, so that there is a preimage $s_U \in \mathcal{F}(NU) = [S\mathcal{F}](U)$ of t_U which then represents a preimage $s_x \in [S\mathcal{F}]_x$ of t_x . This shows that S is also right exact and hence exact.

Now let $\mathcal{F} \xrightarrow{\varphi} \mathcal{G}$ be a morphism of $*$ quasi-coherent \mathcal{O}_{NX} - $*$ modules with $S\varphi = 0$. For faithfulness, we have to show $\varphi = 0$. Choose a finite affine open covering $X = U_1 \cup \dots \cup U_n$ of X . Then $[NX] = [NU_1] \cup \dots \cup [NU_n]$ is a finite $*$ affine $*$ open covering by 4.7, and it suffices to show $\varphi|_{NU_i} = 0$ for all $i \in \{1, \dots, n\}$ or equivalently — because \mathcal{F} and \mathcal{G} are $*$ quasi-coherent — $\varphi_{NU_i} = 0$ for all i . But $\varphi_{NU_i} = [S\varphi]_{U_i} = 0$, and we are done. **q.e.d.**

Let \mathcal{F} and \mathcal{G} be $\mathcal{O}_{N X}$ -*modules. Then N induces a canonical morphism

$$(8) \quad N_* \mathcal{H}om_{\mathcal{O}_{N X}}(\mathcal{F}, \mathcal{G}) \rightarrow \mathcal{H}om_{\mathcal{O}_{\hat{X}}}(N_* \mathcal{F}, N_* \mathcal{G})$$

of $\mathcal{O}_{\hat{X}}$ -modules by

$$\begin{aligned} [N_* \mathcal{H}om_{\mathcal{O}_{N X}}(\mathcal{F}, \mathcal{G})](U) &= \text{Hom}_{\mathcal{O}_{N U}}(\mathcal{F}|_{N U}, \mathcal{G}|_{N U}) \\ &\xrightarrow{N_*} \text{Hom}_{\mathcal{O}_{\hat{U}}}(\mathcal{F}|_U, \mathcal{G}|_U) = [\mathcal{H}om_{\mathcal{O}_{\hat{X}}}(\mathcal{F}, \mathcal{G})](U) \end{aligned}$$

for quasi-compact, open subschemes U of X .

5.9. Proposition. Let \mathcal{F} be a finitely presented \mathcal{O}_X -module, and let \mathcal{G} be an $\mathcal{O}_{N X}$ -*module. Then the canonical morphism (8) (for $N \mathcal{F}$ and \mathcal{G})

$$N_* \mathcal{H}om_{\mathcal{O}_{N X}}(N \mathcal{F}, \mathcal{G}) \longrightarrow \mathcal{H}om_{\mathcal{O}_{\hat{X}}}(N_* N \mathcal{F}, N_* \mathcal{G})$$

of $\mathcal{O}_{\hat{X}}$ -modules is an isomorphism. Taking global sections, this in particular implies that

$$\text{Hom}_{\mathcal{O}_{N X}}(N \mathcal{F}, \mathcal{G}) \xrightarrow{N_*} \text{Hom}_{\mathcal{O}_{\hat{X}}}(N_* N \mathcal{F}, N_* \mathcal{G})$$

is an isomorphism.

Proof. The question whether a given morphism of sheaves on X is an isomorphism is local on X , so we can assume that X is affine. If $\mathcal{F} = \mathcal{O}_X^n$, then $N \mathcal{F} = \mathcal{O}_{N X}^n$ and $N_* N \mathcal{F} = \mathcal{O}_{\hat{X}}^n$, i.e. $\mathcal{H}om_{\mathcal{O}_{N X}}(N \mathcal{F}, \mathcal{G})$ is canonically isomorphic to \mathcal{G}^n (by transfer), and $\mathcal{H}om_{\mathcal{O}_{\hat{X}}}(N_* N \mathcal{F}, N_* \mathcal{G})$ is canonically isomorphic to $N_* \mathcal{G}^n$, so that the statement is obviously true in this case.

In the general case — since X is affine — there is a finite presentation

$$\mathcal{O}_X^m \longrightarrow \mathcal{O}_X^n \longrightarrow \mathcal{F} \longrightarrow 0$$

of \mathcal{F} , which (by 5.3 and 5.4) induces an exact sequence

$$\mathcal{O}_{N X}^m \longrightarrow \mathcal{O}_{N X}^n \longrightarrow N \mathcal{F} \longrightarrow 0$$

of $\mathcal{O}_{N X}$ -*modules and (by 5.8) an exact sequence

$$\mathcal{O}_{\hat{X}}^m \longrightarrow \mathcal{O}_{\hat{X}}^n \longrightarrow N_* N \mathcal{F} \longrightarrow 0$$

of $\mathcal{O}_{\hat{X}}$ -modules. Since the functors

$$\begin{array}{lll} \mathcal{H}om_{\mathcal{O}_{N X}}(-, \mathcal{G}) & : & {}^* \mathcal{M}od_{N X} \rightarrow {}^* \mathcal{M}od_{N X}, \\ \mathcal{H}om_{\mathcal{O}_{\hat{X}}}(-, N_* \mathcal{G}) & : & \mathcal{M}od_{\hat{X}} \rightarrow \mathcal{M}od_{\hat{X}} \quad \text{and} \\ N_* & : & {}^* \mathcal{M}od_{N X} \rightarrow \mathcal{M}od_{\hat{X}} \end{array}$$

are left exact, we get the following commutative diagram of $\mathcal{O}_{\hat{X}}$ -modules with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & N_* \mathcal{H}om_{\mathcal{O}_{N X}}(N \mathcal{F}, \mathcal{G}) & \longrightarrow & N_* \mathcal{H}om_{\mathcal{O}_{N X}}(\mathcal{O}_{N X}^n, \mathcal{G}) & \longrightarrow & N_* \mathcal{H}om_{\mathcal{O}_{N X}}(\mathcal{O}_{N X}^m, \mathcal{G}) \\ & & \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma \\ 0 & \longrightarrow & \mathcal{H}om_{\mathcal{O}_{\hat{X}}}(N_* N \mathcal{F}, N_* \mathcal{G}) & \longrightarrow & \mathcal{H}om_{\mathcal{O}_{\hat{X}}}(\mathcal{O}_{\hat{X}}^n, N_* \mathcal{G}) & \longrightarrow & \mathcal{H}om_{\mathcal{O}_{\hat{X}}}(\mathcal{O}_{\hat{X}}^m, N_* \mathcal{G}). \end{array}$$

According to the first case, β and γ are isomorphisms. But then α must be an isomorphism as well, and we are done. **q.e.d.**

Let \mathcal{F} be a finitely \mathcal{O}_X -module. Choose a subring A_0 of A of finite type over \mathbb{Z} , a scheme X_0 of finite type over A_0 and a finitely presented \mathcal{O}_{X_0} -module \mathcal{F}_0 such that $\langle \mathcal{F}, X/A \rangle$ is the pullback of $\langle \mathcal{F}_0, X_0/A_0 \rangle$ along $\varphi := A_0 \hookrightarrow A$.

By 3.4 and 5.2, we get the following diagram (where we put $\bar{\varphi} := \tau_{\mathbb{Z}, A_0, A}^{-1}[\varphi]$):

$$\begin{array}{ccccc}
 & \mathcal{F} & & N\mathcal{F} & \\
 & \swarrow & & \searrow & \\
 \mathcal{F}_0 & \square & X & \square & {}^*\mathcal{F}_0 \\
 & \swarrow & \searrow f & \swarrow \bar{f} & \searrow \\
 & X_0 & & {}^*X_0 & \\
 & \swarrow & \searrow \varphi^{\text{op}} & \swarrow \bar{\varphi}^{\text{op}} & \searrow \\
 & A_0 & & {}^*A_0 & \\
 & \xrightarrow{*} & & &
 \end{array}$$

The squares are cartesian (in $\mathcal{S}ch^{\text{fp}}$ respectively $\mathcal{M}od^{\text{fp}}$ on the left, in ${}^*\mathcal{S}ch_{*\mathcal{R}}^{\text{fp}}$ respectively ${}^*\mathcal{M}od_{*\mathcal{R}}^{\text{fp}}$ on the right), and we have isomorphisms $f^*\mathcal{F}_0 \xrightarrow{\sim} \mathcal{F}$ and $\bar{f}^*({}^*\mathcal{F}_0) \xrightarrow{\sim} N\mathcal{F}$ and their adjoints $\mathcal{F}_0 \rightarrow f_*\mathcal{F}$ and ${}^*\mathcal{F}_0 \rightarrow \bar{f}_*N\mathcal{F}$.

Now let U_0 be an open subscheme of X_0 , and put $U := U_0 \times_{X_0} X$. We get an $\mathcal{O}_{X_0}(U_0)$ -linear map

$$F_0(U_0) \xrightarrow{*} [{}^*\mathcal{F}_0]({}^*U_0) \longrightarrow [\bar{f}_*N\mathcal{F}]({}^*U_0) = [N\mathcal{F}](NU) = [S N \mathcal{F}](U) = [f_*S N \mathcal{F}](U_0)$$

which is clearly functorial in U_0 , so that we get a morphism of \mathcal{O}_{X_0} -modules $\mathcal{F}_0 \rightarrow f_*S N \mathcal{F}$ and hence — by adjunction — a canonical morphism of \mathcal{O}_X -modules $\mathcal{F} \rightarrow S N \mathcal{F}$.

This morphism is clearly functorial in \mathcal{F} , so that we get a canonical morphism of functors

$$(9) \quad \begin{array}{ccc}
 & \xrightarrow{\text{can}} & \\
 \mathcal{M}od_X^{\text{fp}} & \Downarrow & \mathcal{M}od_X \\
 & \xleftarrow{S N} &
 \end{array}$$

and — again taking adjoints — a canonical morphism of functors

$$(10) \quad \begin{array}{ccc}
 & \xrightarrow{\otimes_{\mathcal{O}_X} \mathcal{O}_{\hat{X}}} & \\
 \mathcal{M}od_X^{\text{fp}} & \Downarrow & \mathcal{M}od_{\hat{X}} \\
 & \xleftarrow{N_* N} &
 \end{array}$$

5.10. Proposition. The canonical morphism of functors (10) is an isomorphism.

Proof. Let \mathcal{F} be a finitely presented \mathcal{O}_X -module. We claim that the canonical morphism $\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_{\hat{X}} \rightarrow N_* N \mathcal{F}$ of $\mathcal{O}_{\hat{X}}$ -modules (or of abelian sheaves on X) is an isomorphism. This claim is local in X , so we can assume that X is affine and choose a finite presentation

$$\mathcal{O}_X^m \rightarrow \mathcal{O}_X^n \rightarrow \mathcal{F} \rightarrow 0.$$

By 5.3, 5.4 and 5.8, we get the following commutative diagram of $\mathcal{O}_{\hat{X}}$ -modules with exact rows:

$$\begin{array}{ccccccc}
 \mathcal{O}_{\hat{X}}^m & \longrightarrow & \mathcal{O}_{\hat{X}}^n & \longrightarrow & \mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_{\hat{X}} & \longrightarrow & 0 \\
 \downarrow \wr & & \downarrow \wr & & \downarrow & & \\
 \mathcal{O}_{\hat{X}}^m & \longrightarrow & \mathcal{O}_{\hat{X}}^n & \longrightarrow & N_* N \mathcal{F} & \longrightarrow & 0.
 \end{array}$$

The first two vertical morphisms are obviously simply the identity, so the third vertical morphism must be an isomorphism. **q.e.d.**

5.11. Proposition. For any affine open subscheme $U = \text{Spec}(B)$ of X , there is a canonical isomorphism of functors

$$(11) \quad \begin{array}{ccc} & \xrightarrow{\Gamma_U(_) \otimes_B N B} & \\ \text{Mod}_X^{\text{fp}} & \Downarrow \wr & [N B]\text{-Mod.} \\ & \xrightarrow{\Gamma_{N U \circ N}} & \end{array}$$

Proof. Using (9), composed with Γ_U , defines a canonical morphism of functors

$$\begin{array}{ccccc} \text{Mod}_X^{\text{fp}} & \xrightarrow{\text{can}} & \text{Mod}_X & \xrightarrow{\Gamma_U} & B\text{-Mod} \\ & \searrow N & \downarrow (9) & \nearrow S & \\ & & {}^*\text{Mod}_{N X}^{\text{fp}} & \xrightarrow{\Gamma_{N U}} & [N B\text{-Mod}] \end{array} \quad \begin{array}{c} \\ \\ = \\ \nearrow \text{can} \end{array}$$

and thus by adjunction the morphism of functors (11). To see that this is an isomorphism, let \mathcal{F} be a finitely presented \mathcal{O}_X -module, and choose a finite presentation

$$B^m \longrightarrow B^n \longrightarrow \mathcal{F}(U) \longrightarrow 0.$$

Taking associated sheaves and applying N , we get an exact sequence of * finitely presented $\mathcal{O}_{N U}$ - * modules

$$\mathcal{O}_{N U}^m \longrightarrow \mathcal{O}_{N U}^n \longrightarrow [N \mathcal{F}]|_{N U} \longrightarrow 0.$$

By transfer, $\Gamma_{N U} : \text{Mod}_{N U}^{\text{fp}} \longrightarrow [N B]\text{-Mod}$ is exact, so we get the exact sequence of $N B$ -modules

$$N B^m \longrightarrow N B^n \longrightarrow [N \mathcal{F}](N U) \longrightarrow 0$$

and hence the following commutative diagram of $N B$ -modules with exact rows:

$$\begin{array}{ccccccc} B^m \otimes_B N B & \longrightarrow & B^n \otimes_B N B & \longrightarrow & \mathcal{F}(U) \otimes_B N B & \longrightarrow & 0 \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \\ N B^m & \longrightarrow & N B^n & \longrightarrow & [N \mathcal{F}](N U) & \longrightarrow & 0. \end{array}$$

Since α and β are clearly isomorphisms, so is γ , and we are done. **q.e.d.**

5.12. Corollary. The canonical functors

$$\begin{array}{ccc} (\text{Mod}_X^{\text{fp}})^{\text{op}} \times {}^*\text{Mod}_{N X} & \longrightarrow & \text{Sets} \\ (\mathcal{F} \quad , \quad \mathcal{G}) & \mapsto & \begin{cases} \text{Hom}_{\mathcal{O}_{N X}}(N \mathcal{F}, \mathcal{G}) \\ \text{Hom}_{\mathcal{O}_X}(\mathcal{F}, S \mathcal{G}) \end{cases} \end{array}$$

are canonically isomorphic via

$$\tau_{X, \mathcal{F}, \mathcal{G}} : \text{Hom}_{\mathcal{O}_{N X}}(N \mathcal{F}, \mathcal{G}) \longrightarrow \text{Hom}_{\mathcal{O}_X}(\mathcal{F}, S \mathcal{G}), \quad [N \mathcal{F} \xrightarrow{\varphi} \mathcal{G}] \mapsto [\mathcal{F} \xrightarrow{(9)} S N \mathcal{F} \xrightarrow{S \varphi} S \mathcal{G}].$$

Proof. Let \mathcal{F} be a finitely presented \mathcal{O}_X -module, and let \mathcal{G} be an $\mathcal{O}_{N X}$ - * module. Then

$$\text{Hom}_{\mathcal{O}_{N X}}(N \mathcal{F}, \mathcal{G}) \stackrel{5.9}{\cong} \text{Hom}_{\mathcal{O}_{\hat{X}}}(N {}^*\mathcal{F}, N {}^*\mathcal{G}) \stackrel{5.10}{\cong} \text{Hom}_{\mathcal{O}_{\hat{X}}}(\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_{\hat{X}}, N {}^*\mathcal{G}) \stackrel{\text{adj.}}{\cong} \text{Hom}_{\mathcal{O}_X}(\mathcal{F}, S \mathcal{G}),$$

and it is clear that the composition of these canonical isomorphisms is just $\tau_{X, \mathcal{F}, \mathcal{G}}$. **q.e.d.**

5.13. Remark. Note that the functoriality of the isomorphism from 5.12 in particular implies the following: If $\mathcal{F} \xrightarrow{\varphi} \mathcal{F}'$ is a morphism of finitely presented \mathcal{O}_X -modules, if $\mathcal{G} \xrightarrow{\psi} \mathcal{G}'$ is a morphism of \mathcal{O}_{NX} -*modules, and if

$$(12) \quad \begin{array}{ccc} N\mathcal{F} & \xrightarrow{N\varphi} & N\mathcal{F}' \\ f \downarrow & & \downarrow g \\ \mathcal{G} & \xrightarrow{\psi} & \mathcal{G}' \end{array}$$

is a diagram of \mathcal{O}_{NX} -*modules, then (12) commutes if and only if the corresponding diagram

$$(13) \quad \begin{array}{ccc} \mathcal{F} & \xrightarrow{\varphi} & \mathcal{F}' \\ \tau_{X,\mathcal{F},\mathcal{G}}(f) \downarrow & & \downarrow \tau_{X,\mathcal{F}',\mathcal{G}'}(g) \\ S\mathcal{G} & \xrightarrow{S\psi} & S\mathcal{G}' \end{array}$$

of \mathcal{O}_X -modules commutes.

5.14. Corollary. Let \mathcal{F} and \mathcal{G} be two finitely presented \mathcal{O}_X -modules. There is a canonical isomorphism of *finitely presented \mathcal{O}_{NX} -modules

$$N(\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}) \xrightarrow{\sim} N\mathcal{F} \otimes_{\mathcal{O}_{NX}} N\mathcal{G}.$$

Proof. For a quasi-compact open subscheme U of X , we have a canonical $\mathcal{O}_X(U)$ -linear map

$$\begin{aligned} \mathcal{F}(U) \otimes_{\mathcal{O}_X(U)} \mathcal{G}(U) &\xrightarrow{(9)} [S N \mathcal{F}](U) \otimes_{[S N \mathcal{O}_X](U)} [S N \mathcal{G}](U) \\ &= [N \mathcal{F}](NU) \otimes_{\mathcal{O}_{NX}(NU)} [N \mathcal{G}](NU) \xrightarrow{\text{can}} [N \mathcal{F} \otimes_{\mathcal{O}_{NX}} N \mathcal{G}](NU) = S[N \mathcal{F} \otimes_{\mathcal{O}_{NX}} N \mathcal{G}](U), \end{aligned}$$

which is clearly functorial in U and consequently defines a functorial morphism of *presheaves* of \mathcal{O}_X -modules

$$[U \mapsto \mathcal{F}(U) \otimes_{\mathcal{O}_X(U)} \mathcal{G}(U)] \longrightarrow S[N \mathcal{F} \otimes_{\mathcal{O}_{NX}} N \mathcal{G}]$$

and then, by the universal property of the associated sheaf, a functorial morphism of \mathcal{O}_X -modules

$$\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G} \longrightarrow S[N \mathcal{F} \otimes_{\mathcal{O}_{NX}} N \mathcal{G}],$$

which by 5.12 and 5.13 corresponds to a functorial morphism of \mathcal{O}_{NX} -*modules

$$(14) \quad N[\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}] \longrightarrow N\mathcal{F} \otimes_{\mathcal{O}_{NX}} N\mathcal{G}.$$

To prove that (14) is an isomorphism, choose a quadruple $\langle A_0, X_0, \mathcal{F}_0, \mathcal{G}_0 \rangle$, where $A_0 \xrightarrow{\varphi} A$ is a finitely generated subring of A , X_0 is an A_0 -scheme of finite type with $X \cong \varphi^* X_0$ and \mathcal{F}_0 and \mathcal{G}_0 are coherent sheaves on X_0 with $\mathcal{F} \cong \varphi^* \mathcal{F}_0$ and $\mathcal{G} \cong \varphi^* \mathcal{G}_0$. Then of course we also have $\varphi^*[\mathcal{F}_0 \otimes_{\mathcal{O}_{X_0}} \mathcal{G}_0] \cong \mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}$ and therefore (with $\bar{\varphi} := \tau_{\mathbb{Z}, A_0, A}^{-1}[\varphi]$)

$$\begin{aligned} N[\mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{G}] &\cong \bar{\varphi}^* \left(*[\mathcal{F}_0 \otimes_{\mathcal{O}_{X_0}} \mathcal{G}_0] \right) = \bar{\varphi}^* \left(*\mathcal{F}_0 \otimes_{\mathcal{O}_{*X_0}} *\mathcal{G}_0 \right) \\ &\cong \bar{\varphi}^*(*\mathcal{F}_0) \otimes_{\mathcal{O}_{NX}} \bar{\varphi}^*(*\mathcal{G}_0) \cong N\mathcal{F} \otimes_{\mathcal{O}_{NX}} N\mathcal{G} \end{aligned}$$

by construction of N . **q.e.d.**

5.15. Corollary. The functor $N : \mathcal{M}od_X^{\text{fp}} \longrightarrow *\mathcal{M}od_{NX}^{\text{fp}}$ induces a canonical group homomorphism $N : \text{Pic}(X) \longrightarrow *\text{Pic}(NX)$ between the Picard group of X and the *Picard group of NX .

Proof. By 5.5, N sends line bundles to line bundles, so we get a map $N : \text{Pic}(X) \longrightarrow *\text{Pic}(NX)$. This map is a group homomorphism by 5.14. **q.e.d.**

5.16. Corollary. and let \mathcal{F} and \mathcal{G} be two finitely presented \mathcal{O}_X -modules with the property that the \mathcal{O}_X -module $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})$ is also finitely-presented, which is for example the case if

- \mathcal{F} is a vector bundle or
- \mathcal{F} and \mathcal{G} are coherent.

Then there is a canonical morphism of $*$ finitely presented $\mathcal{O}_{N X}$ - $*$ modules

$$(15) \quad N \mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) \longrightarrow \mathcal{H}om_{\mathcal{O}_{N X}}(N \mathcal{F}, N \mathcal{G})$$

which is an isomorphism if \mathcal{F} is a vector bundle.

Proof. Look at the following canonical map of sets of morphisms:

$$\begin{aligned} \text{Mor}_{\mathcal{M}od_X^{\text{fp}}}(\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}), \mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})) &\cong \text{Mor}_{\mathcal{M}od_X^{\text{fp}}}(\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) \otimes_{\mathcal{O}_X} \mathcal{F}, \mathcal{G}) \\ &\xrightarrow{N} \text{Mor}_{*\mathcal{M}od_{N X}^{\text{fp}}}(N[\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) \otimes_{\mathcal{O}_X} \mathcal{F}], N \mathcal{G}) \\ &\stackrel{5.14}{\cong} \text{Mor}_{*\mathcal{M}od_{N X}^{\text{fp}}}(N \mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) \otimes_{\mathcal{O}_{N X}} N \mathcal{F}, N \mathcal{G}) \\ &\cong \text{Mor}_{*\mathcal{M}od_{N X}^{\text{fp}}}(N \mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}), \mathcal{H}om_{\mathcal{O}_{N X}}(N \mathcal{F}, N \mathcal{G})), \end{aligned}$$

and take the identity's image under this map to get (15).

Now let \mathcal{F} be a vector bundle. Since the question whether (15) is an isomorphism is local, we can assume that $\mathcal{F} = \mathcal{O}_X^n$ is trivial, and we have

$$N \mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) \cong N \mathcal{G}^n \stackrel{\text{transfer}}{\cong} \text{Hom}_{N X}(\mathcal{O}_{N X}^n, N \mathcal{G}) \stackrel{5.4}{\cong} \text{Hom}_{N X}(N \mathcal{O}_X^n, N \mathcal{G})$$

as desired. **q.e.d.**

5.17. Corollary. For a vector bundle \mathcal{E} on X , there is a canonical isomorphism $N(\mathcal{E}^\vee) \cong (N \mathcal{E})^\vee$.

Proof. This follows immediately from 5.4 and 5.16:

$$N(\mathcal{E}^\vee) = N \mathcal{H}om_{\mathcal{O}_X}(\mathcal{E}, \mathcal{O}_X) \stackrel{5.16}{\cong} \mathcal{H}om_{\mathcal{O}_{N X}}(N \mathcal{E}, N \mathcal{O}_X) \stackrel{5.4}{\cong} \mathcal{H}om_{\mathcal{O}_{N X}}(N \mathcal{E}, \mathcal{O}_{N X}) = (N \mathcal{E})^\vee.$$

q.e.d.

5.18. Proposition. Let $n \in \mathbb{N}_+$ be a natural number, and let $k \in \mathbb{Z}$ be an integer.

- Under the functor N , the invertible $\mathcal{O}_{\mathbb{P}_X^n}$ -module $\mathcal{O}_{\mathbb{P}_X^n}(k)$ is mapped to the $*$ invertible $\mathcal{O}_{*\mathbb{P}_{N X}^n}$ -module $\mathcal{O}_{*\mathbb{P}_{N X}^n}(k)$.
- Let $i : Y \hookrightarrow \mathbb{P}_X^n$ be a closed immersion of finitely presented A -schemes, and let \mathcal{F} be a finitely presented \mathcal{O}_Y -module. Then $N[\mathcal{F}(k)] = [N \mathcal{F}](k)$, where the twists are taken with respect to i respectively $N i$.

Proof. Choose a finitely generated subring $A_0 \xrightarrow{\varphi} A$ of A and an A_0 -scheme X_0 of finite type with $\varphi^* X_0 = X$. Then $\varphi^* \mathbb{P}_{X_0}^n = \mathbb{P}_X^n$, so

$$N(\mathcal{O}_{\mathbb{P}_X^n}(k)) = \left[\tau_{\mathbb{Z}, A_0, A}^{-1}[\varphi] \right]^* \mathcal{O}_{*\mathbb{P}_{X_0}^n}(k) = \mathcal{O}_{N(\mathbb{P}_X^n)}(k) \stackrel{3.7}{\cong} \mathcal{O}_{*\mathbb{P}_{N X}^n}(k),$$

and this is $*$ invertible by 5.5 (or by transfer), so we have (i).

For (ii) we get:

$$\begin{aligned} N[\mathcal{F}(k)] &= N[\mathcal{F} \otimes_{\mathcal{O}_Y} i^* \mathcal{O}_{\mathbb{P}_X^n}(k)] \stackrel{5.14}{\cong} [N \mathcal{F}]^* \otimes_{\mathcal{O}_{N Y}} [N[i^* \mathcal{O}_{\mathbb{P}_X^n}(k)]] \\ &= [N \mathcal{F}]^* \otimes_{\mathcal{O}_{N Y}} [(N i)^* [N \mathcal{O}_{\mathbb{P}_X^n}(k)]] \stackrel{(i)}{\cong} [N \mathcal{F}]^* \otimes_{\mathcal{O}_{N Y}} [(N i)^* [\mathcal{O}_{*\mathbb{P}_{N X}^n}(k)]] = [N \mathcal{F}](k). \end{aligned}$$

q.e.d.

If Z is a finitely presented closed subscheme of X , given by a finitely presented sheaf of ideals I on X , then we know from 4.4 that NZ is a $*$ closed $*$ subscheme of NX . As final result in this section, we want to determine the relationship between NI and the $*$ ideal on NX defining NZ :

5.19. Proposition. Let Z be a finitely presented closed subscheme of X , given by a finitely presented sheaf of ideals I . Then the $*$ closed $*$ subscheme NZ of NX is given by the $*$ ideal $\text{Im}(NI \rightarrow \mathcal{O}_{NX})$.

Proof. Let NZ be given by the $*$ ideal \mathcal{J} on NX . If $U \subseteq X$ is a quasi-compact open subscheme of X , then the $*$ closed $*$ subscheme $N[Z \cap U]$ of NU is given by $\mathcal{J}|_{NU}$, so we can assume without loss of generality that X is affine, say $X = \text{Spec}(B)$ for a finitely presented A -algebra B . Then $Z = \text{Spec}(B/\mathfrak{b})$ for a finitely presented ideal \mathfrak{b} of B , and $I = \tilde{\mathfrak{b}}$.

Using 4.15, we have

$$N[B/\mathfrak{b}] = [NB]/\mathfrak{b} = [NB]/\mathfrak{b} \cdot [NB],$$

so that NZ is given by the $*$ ideal $\mathcal{J} = \widetilde{\mathfrak{b} \cdot [NB]}$ of NX , and we have to prove that this $*$ ideal equals $\text{Im}(NI \rightarrow \mathcal{O}_{NX})$ or — equivalently — that the global sections of these two $*$ ideals agree (as ideals of NB). Using 5.11, this is easy:

$$\Gamma_{NX}[\text{Im}(NI \rightarrow \mathcal{O}_{NX})] = \text{Im}(\Gamma_{NX}[NI] \rightarrow NB) \stackrel{(11)}{=} \text{Im}(\mathfrak{b} \otimes_B [NB] \rightarrow NB) = \mathfrak{b} \cdot NB = \Gamma_{NX}[\mathcal{J}].$$

q.e.d.

6. THE CASE OF VARIETIES

Let k be a $*$ field in $*$ \mathcal{R} , i.e. a $*$ ring which is an (internal) field. Then k is of course a noetherian ring, so that a k -scheme X is finitely presented if and only if it is of finite type, and an \mathcal{O}_X -module \mathcal{F} is finitely presented if and only if it is coherent.

6.1. Definition. We can consider "dimension" as a function $\dim : \{\text{schemes}\} \longrightarrow \{-\infty\} \amalg \mathbb{N}_0 \amalg \{\infty\}$, so by restriction to $\text{Ob}(\mathcal{S})$ and enlarging we get an induced function

$$*\dim : \{*\text{schemes}\} \longrightarrow \{-\infty\} \amalg *\mathbb{N}_0 \amalg \{\infty\}.$$

For a $*$ scheme X , we call $*\dim X$ the *$*$ dimension of X* .

For the proof of theorem 6.4 below, we will need the following results of van den Dries and Schmidt which we state here — in our notation — for the convenience of the reader:

6.2. Theorem. (Lou van den Dries, K. Schmidt)

Let $I \subseteq k[X_1, \dots, X_n]$ be an ideal. Then

- (i) The ring homomorphism $k[X_i] \rightarrow k^*[X_i]$ is faithfully flat.
- (ii) I is prime if and only if $*I \subseteq k^*[X_i]$ is $*$ prime or — what amounts to the same, since for an ideal being prime is clearly a first order property — prime.
- (iii) If $\mathfrak{p}_1, \dots, \mathfrak{p}_m$ are the distinct minimal primes of I , then $*\mathfrak{p}_1, \dots, *\mathfrak{p}_m$ are the distinct minimal primes of $*I$ (in particular, all minimal primes of $*I$ are $*$ ideals, hence the notions of "minimal prime ideal of $*I$ " and "minimal prime $*$ ideal of $*I$ " coincide).
- (iv) $\sqrt[*]{*I} = *[\sqrt{I}]$.

Proof. Part (i) is [vdDS84, 1.8], part (ii) is [vdDS84, 2.5], and parts (iii) and (iv) are [vdDS84, 2.7]; that van den Dries and Schmidt's formulation agrees with the one given here follows immediately from 2.8.
q.e.d.

6.3. Corollary. Let A be a k -algebra of finite type. Then $\sigma_A : A \rightarrow NA$ is faithfully flat.

Proof. Let $A = k[X_1, \dots, X_N]/I$. Then

$$\sigma_A = \sigma_{k[X_i]} \otimes_{k[X_i]} k[X_i]/I \quad : \quad A = k[X_i]/I \longrightarrow k^*[X_i]/I \cdot k^*[X_i] \stackrel{2.8}{=} k^*[X_i]/^*I = N A,$$

and $\sigma_{k[X_i]}$ is faithfully flat by 6.2(i), so σ_A — as a base change of $\sigma_{k[X_i]}$ — must be faithfully flat as well. **q.e.d.**

6.4. Theorem. Let X be a scheme of finite type over k .

- (i) X is the empty scheme if and only if $N X$ is the * empty scheme.
- (ii) $^* \dim N X = \dim X$.
- (iii) X is reduced (irreducible, integer) if and only if $N X$ is * reduced (* irreducible, * integer).
- (iv) The functor N from coherent \mathcal{O}_X -modules to (the abelian category of) * coherent $\mathcal{O}_{N X}$ -modules is faithful and exact.

Proof. If $X = \emptyset$, then $N X = ^*\emptyset$ by 4.1(iii), so let $N X = ^*\emptyset$. Let us assume that $X \neq \emptyset$. Then X contains a K -valued point for a finite field extension K/k , and applying N gives us an $N K$ -valued point of $N X$. If $K = k[X_1, \dots, X_n]/I$ is a finite presentation of K , it follows from 6.2(i) that $N K = k^*[X_i]/^*I$ is not zero, so the existence of an $N K$ -valued point of $N X$ proves the existence of a * topological point of $N X$, a contradiction to $N X = ^*\emptyset$.

Having settled (i), for (ii) and (iii) we can assume that $X \neq \emptyset$. For (ii), we use [EGA IV₂, 4.1.2], according to which $\dim X = n$ is equivalent to the existence of a diagram

$$\begin{array}{ccc} U & \xrightarrow{j} & X \\ f \downarrow & & \\ \mathbb{A}_k^n & & \end{array}$$

of k -schemes of finite type with an open immersion j and a finite and surjective f . But then (ii) follows from 4.4 and from the transfer of [EGA IV₂, 4.1.2].

For (iii), note that we only have to prove the claim for "reduced" and "irreducible", since "integer" is just the conjunction of those two.

Let us first consider the case where $X = \operatorname{Spec}(k[X_1, \dots, X_n]/I)$ is affine. We have

$$\begin{aligned} X \text{ reduced} &\iff \sqrt{I} = I \iff \sqrt{I}/I = (0) \stackrel{6.2(i)}{\iff} \sqrt{I} \cdot k^*[X_i]/I \cdot k^*[X_i] = (0) \\ &\stackrel{2.8}{\iff} ^*[\sqrt{I}] = ^*I \stackrel{6.2(iv)}{\iff} \sqrt{^*I} = ^*I \iff N X \text{ } ^*\text{reduced} \end{aligned}$$

and

X irreducible $\iff I$ has exactly one minimal prime ideal

$$\stackrel{6.2(iii)}{\iff} ^*I \text{ has exactly one minimal prime } ^*\text{ideal} \iff N X \text{ } ^*\text{irreducible}.$$

In the general case, let $(U_j)_{j \in J}$ be a finite open covering of X by affine schemes U_j which are not empty. The scheme X is reduced if and only if the U_j are reduced, which we have just proven to be equivalent to the $N U_j$ being * reduced, which in turn is equivalent to $N X$ being * reduced by 4.7 and transfer.

Let X be irreducible. Then all U_j are irreducible, and their intersection is an open non-empty subscheme of X . Then by (i), the * scheme

$$N \bigcap_{j \in J} U_j \stackrel{4.1(i)}{=} \bigcap_{j \in J} N U_j$$

is not * empty. Since we already know that the $N U_j$ are * irreducible and therefore * connected, this implies that $N X$ is * connected.

Assume that $N X$ is * reduced. Since $N X$ is * connected, there must be a * topological point of $N X$ where two * irreducible components of $N X$ intersect, and since the $N U_j$ cover $N X$, this * topological point lies in one of the U_j which consequently can not be * irreducible, a contradiction.

Now let $N X$ be * irreducible, and assume that X is not irreducible. Since $N X$ is * irreducible, the $N U$ are * irreducible, and their intersection is not * empty, so by (i) and 4.1(i), the scheme X is connected.

Reasoning as above, we see this implies that one of the U_j is reducible, which contradicts the fact that the NU_j are \ast irreducible.

For (iv), we have to show that a short sequence of coherent \mathcal{O}_X -modules

$$0 \longrightarrow \mathcal{F}' \longrightarrow \mathcal{F} \longrightarrow \mathcal{F}'' \longrightarrow 0$$

is exact if and only if the induced sequence of \ast coherent \mathcal{O}_{NX} -modules

$$0 \longrightarrow N\mathcal{F}' \longrightarrow N\mathcal{F} \longrightarrow N\mathcal{F}'' \longrightarrow 0$$

is exact, which by 5.8 and 5.10 is equivalent to the exactness of

$$0 \longrightarrow \mathcal{F}' \otimes_{\mathcal{O}_X} \mathcal{O}_{\hat{X}} \longrightarrow \mathcal{F} \otimes_{\mathcal{O}_X} \mathcal{O}_{\hat{X}} \longrightarrow \mathcal{F}'' \otimes_{\mathcal{O}_X} \mathcal{O}_{\hat{X}} \longrightarrow 0.$$

Taking stalks, it is enough to show that for every point $x \in X$,

$$0 \longrightarrow \mathcal{F}'_x \longrightarrow \mathcal{F}_x \longrightarrow \mathcal{F}''_x \longrightarrow 0$$

is exact if and only if

$$0 \longrightarrow \mathcal{F}'_x \otimes_{\mathcal{O}_{X,x}} \mathcal{O}_{\hat{X},x} \longrightarrow \mathcal{F}_x \otimes_{\mathcal{O}_{X,x}} \mathcal{O}_{\hat{X},x} \longrightarrow \mathcal{F}''_x \otimes_{\mathcal{O}_{X,x}} \mathcal{O}_{\hat{X},x} \longrightarrow 0$$

is exact. But since

$$\mathcal{O}_{\hat{X},x} = \varinjlim_{x \in U \subseteq X} \mathcal{O}_{\hat{X}}(U) = \varinjlim_{x \in U \subseteq X} N[\mathcal{O}_X(U)],$$

where the limit is taken over all affine neighborhoods of x in X , we see from 6.3 that $\mathcal{O}_{X,x} \longrightarrow \mathcal{O}_{\hat{X},x}$ is faithfully flat, and the claim follows. **q.e.d.**

6.5. Corollary. Let X be a k -scheme of finite type, and let \mathcal{F} and \mathcal{G} be coherent \mathcal{O}_X -modules. Then the canonical morphism (15) is an isomorphism:

$$N\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) \xrightarrow{\sim} \mathcal{H}om_{\mathcal{O}_{NX}}(N\mathcal{F}, N\mathcal{G}).$$

Proof. Since the question is local, we can assume that there exists a global presentation

$$\mathcal{O}_X^m \longrightarrow \mathcal{O}_X^n \longrightarrow \mathcal{F} \longrightarrow 0,$$

and since N is exact by 6.4(iv), the functors $N\mathcal{H}om_{\mathcal{O}_X}(_, \mathcal{G})$ and $\mathcal{H}om_{\mathcal{O}_{NX}}(N_, N\mathcal{G})$ from \mathcal{Coh}_X to $\ast\mathcal{Coh}_{NX}$ are both left exact, so that we get the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & N\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) & \longrightarrow & N\mathcal{H}om_{\mathcal{O}_X}(\mathcal{O}_X^n, \mathcal{G}) & \longrightarrow & N\mathcal{H}om_{\mathcal{O}_X}(\mathcal{O}_X^m, \mathcal{G}) \\ & & \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma \\ 0 & \longrightarrow & \mathcal{H}om_{\mathcal{O}_{NX}}(N\mathcal{F}, N\mathcal{G}) & \longrightarrow & \mathcal{H}om_{\mathcal{O}_{NX}}(N\mathcal{O}_X^n, N\mathcal{G}) & \longrightarrow & \mathcal{H}om_{\mathcal{O}_{NX}}(N\mathcal{O}_X^m, N\mathcal{G}) \end{array}$$

with the vertical morphisms given by (15). By 5.16, both β and γ are isomorphisms, so α must be an isomorphism as well. **q.e.d.**

6.6. Corollary. Let X be a k -scheme of finite type, and let Z be a closed subscheme of X corresponding to a sheaf of ideals I on X . Then the \ast closed \ast subscheme NZ of NX is given by the \ast ideal NI .

Proof. According to 5.19, NZ is given by the \ast ideal $\text{Im}(NI \rightarrow \mathcal{O}_{NX})$. But $I \rightarrow \mathcal{O}_X$ is a monomorphism and N is exact by 6.4(iv), so $NI \hookrightarrow \mathcal{O}_{NX}$, and the corollary follows. **q.e.d.**

6.7. Lemma. Let A be a finitely generated k -algebra, let I be an ideal of A , and let $f \in A$. Consider the ideals $(I : f^n) := \{a \in A \mid af^n \in I\}$ (for $n \in \mathbb{N}_+$) and $(I : f^\infty) := \bigcup_{n \in \mathbb{N}_+} (I : f^n)$ of A . Then

$$(16) \quad \forall n \in \mathbb{N}_+ : (I : f^n) \cdot NA = (I \cdot NA : f^n)$$

and

$$(17) \quad (I : f^\infty) \cdot NA = \bigcup_{n \in \ast\mathbb{N}_+} (I \cdot NA : f^n) = (I \cdot NA : f^\infty).$$

Proof. By definition, the diagram

$$\begin{array}{ccc} (I : f^n) & \longrightarrow & I \\ \downarrow & \square & \downarrow \\ A & \xrightarrow{f^n} & A \end{array}$$

is cartesian in the category of A -modules for every $n \in \mathbb{N}_+$. Since $A \xrightarrow{\sigma_A} N A$ is (faithfully) flat by 6.3, this implies (16). For (17) note that $(I : f^\infty)$ is finitely generated, because A is noetherian. Consequently, there is an $N \in \mathbb{N}_+$ with $(I : f^N) = (I : f^{N+1})$ and hence $(I : f^n) = (I : f^N)$ for all $n \geq N$ and $(I : f^\infty) = (I : f^N)$. Then

$$(I \cdot N A : f^N) \stackrel{(16)}{=} (I : f^N) \cdot N A = (I : f^{N+1}) \cdot N A \stackrel{(16)}{=} (I \cdot N A : f^{N+1}),$$

and hence $(I \cdot N A : f^n) = (I \cdot N A : f^N)$ for all $n \geq N$ by transfer — so (17) holds. **q.e.d.**

6.8. Proposition. Let X be a k -scheme of finite type, let $Y \subseteq X$ be a subscheme, and let $\bar{Y} \subseteq X$ be the scheme theoretic closure of Y in X . Then $N \bar{Y}$ is the *scheme theoretic closure of $N Y$ in $N X$.

Proof. If $U \subseteq X$ is an open subscheme, then $\overline{Y \cap U}$, the closure of $Y \cap U$ in U , equals $\bar{Y} \cap U$. Therefore we can assume without loss of generality that $X = \text{Spec}(A)$ is affine and that $Y = \text{Spec}(A/I) \cap D(f_1) \cap \dots \cap D(f_n)$ for an ideal $I \subseteq A$ and elements $f_1, \dots, f_n \in A$. Then $\bar{Y} = \text{Spec}(A/\mathfrak{J})$ with $\mathfrak{J} = \bigcap_{i=1}^n \ker[A \xrightarrow{\text{can}} A_{f_i}/I A_{f_i}]$. For any $f \in A$, we have

$$\ker[A \rightarrow A_f/I A_f] = \{a \in A \mid \exists n \in \mathbb{N}_+ : f^n a \in I\} = \bigcup_{n=1}^{\infty} (I : f^n) = (I : f^\infty),$$

so $\mathfrak{J} = \bigcup_{i=1}^n (I : f_i^\infty)$. Let $\tilde{\mathfrak{J}} \subseteq N A$ be the *ideal corresponding to the *scheme theoretic closure of $N Y$ in $N X$. By transfer, we have

$$\tilde{\mathfrak{J}} = \bigcap_{i=1}^n (I \cdot N A : f_i^\infty) \stackrel{6.7}{=} \bigcap_{i=1}^n (I : f_i^\infty) \cdot N A = \mathfrak{J} \cdot N A,$$

so $[N A]/\tilde{\mathfrak{J}} = N[A/\mathfrak{J}]$, and we are done. **q.e.d.**

6.9. Proposition. Let X be a k -scheme of finite type, let $Y \subseteq X$ be a closed subscheme, and let $f : Z \rightarrow X$ be the blow-up of X in Y . Then $N f : N Z \rightarrow N X$ is the *blow-up of $N X$ in $N Y$.

Proof. First note that $N Y$ is a *closed *subscheme of $N X$ by 4.4, so the statement makes sense. Next, by [EGA IV₃, 8.8.2, 8.10.5] there exist a finitely generated subring k_0 of k , a k_0 -scheme X_0 of finite type with $X = X_0 \times_{k_0} k$ and a closed subscheme Y_0 of X_0 with $Y = Y_0 \times_{k_0} k$.

Let $Z_0 \rightarrow X_0$ be the blow-up of X_0 in Y_0 , and let W be the cartesian diagram of k -schemes

$$(18) \quad \begin{array}{ccccc} W & \xrightarrow{\quad \hookrightarrow \quad} & X \times_{X_0} Z_0 & \longrightarrow & Z_0 \\ \downarrow & \square & \downarrow \pi & \square & \downarrow \pi_0 \\ X \setminus Y & \xrightarrow{\quad \hookrightarrow \quad} & X & \longrightarrow & X_0. \end{array}$$

If $Z := \bar{W}$ denotes the scheme theoretic closure of W in $X \times_{X_0} Z_0$, then $f := \pi|_Z : Z \rightarrow X$ is the blow-up of X in Y (compare [EH00, IV-21]).

Applying the functor N to the left square of (18) and using 4.1(i) and 4.8, we get a cartesian square of k -*schemes

$$\begin{array}{ccc} N W & \xrightarrow{\quad \hookrightarrow \quad} & [N X] \times_{*_{X_0}} * Z_0 \\ \downarrow & \square & \downarrow N \pi \\ [N X] \setminus [N Y] & \xrightarrow{\quad \hookrightarrow \quad} & N X, \end{array}$$

and by transfer, the *blow-up of $N X$ in $N Y$ is the *scheme theoretic closure of $N W$ in $[N X] \times_{*_{X_0}} * Z_0$. But according to 6.8, this is just $N \bar{W} = N Z$, which completes the proof. **q.e.d.**

6.10. Definition. For every field K , every K -scheme X and every K -rational point $x \in X$, we have the K -vector space $T_{X,x}$, the *(Zariski) tangent space of X at x* , defined as the K -dual of $\mathfrak{m}_x/\mathfrak{m}_x^2$.

By transfer, for every $*$ field K , every $*$ scheme X over K and every K -valued point x of X , we thus have an internal K -vector space $T_{X,x}$ which we also call the *(Zariski) tangent space of X at x* .

6.11. Proposition. let X be a k -scheme of finite type, and let $x \in X$ be a k -rational point. Then N induces a canonical functorial k -isomorphism of Zariski tangent spaces

$$N : T_{X,x} \xrightarrow{\sim} T_{NX, Nx}.$$

Proof. Identify x with a k -morphism $x : \text{Spec}(k) \rightarrow X$, and let $e : \text{Spec}(k) \rightarrow \text{Spec}(k[\varepsilon]/\varepsilon^2)$ be the k -morphism induced by sending ε to zero.

It is well known that there is a canonical functorial isomorphism of k -vector spaces

$$(19) \quad T_{X,x} \cong \{t \in X(k[\varepsilon]/\varepsilon^2) \mid e^*t = x\}.$$

By transfer, we get a canonical functorial isomorphism of internal k -vector spaces

$$(20) \quad T_{NX, Nx} \cong \{t \in (NX)(k^*[\varepsilon]/\varepsilon^2) \mid (Ne)^*t = Nx\}.$$

But by 4.16 we have $k^*[\varepsilon]/\varepsilon^2 = k[\varepsilon]/\varepsilon^2$, and we get the following commutative diagram of sets:

$$\begin{array}{ccc} X(k[\varepsilon]/\varepsilon^2) & \xrightarrow{e^*} & X(k) \\ N \downarrow \wr & & \downarrow \wr N \\ (NX)(k[\varepsilon]/\varepsilon^2) & \xrightarrow{(Ne)^*} & (NX)(k), \end{array}$$

where the vertical maps are bijections because of 4.13. From this, (19) and (20) the claim immediately follows. **q.e.d.**

6.12. Corollary. Assume that k is $*$ algebraically closed, and let X be a k -scheme of finite type. If NX is $*$ nonsingular, then X is nonsingular.

Proof. Let $d := \dim X$, and let $x \in X$ be a closed point. Since k is $*$ algebraically closed, k is externally an algebraically closed field, and x is a k -rational point. Since NX is $*$ nonsingular of $*$ dimension d (by 6.4(ii)), the tangent space $T_{NX, Nx}$ has $*$ dimension d , and the tangent space $T_{X,x}$ has dimension d by 6.11. This shows that all tangent spaces of X at closed points have dimension d , which means that X is nonsingular. **q.e.d.**

6.13. Proposition. Let X be a k -scheme of finite type, and let Y and Z be two subschemes of X . If $NY \hookrightarrow NX$ factors through $NZ \hookrightarrow NX$, then $Y \hookrightarrow X$ factors through $Z \hookrightarrow X$. In particular, if NY and NZ are the same $*$ subspaces of NX , then Y and Z are the same subschemes of X .

Proof. Factor $Z \hookrightarrow X$ as $Z \xrightarrow{i_Z} V \xrightarrow{j_Z} X$ with a closed immersion i_Z and an open immersion j_Z . We claim that $Y \hookrightarrow X$ factors through j_Z : Equip $Y \setminus V$ with its reduced structure and consider the cartesian diagram

$$\begin{array}{ccc} \emptyset & \hookrightarrow & Y \setminus V \\ \downarrow & \square & \downarrow \\ V & \xrightarrow{j} & X. \end{array}$$

Applying N and using 4.1(i) and 4.1(iii), we get a cartesian diagram

$$(21) \quad \begin{array}{ccc} \emptyset^c & \xrightarrow{\quad} & N[Y \setminus V] \\ \downarrow & \square & \downarrow \\ NV^c & \xrightarrow{Nj} & NX. \end{array}$$

If $Y \setminus V$ was not empty, then $N[Y \setminus V]$ also would not be empty by 6.4(i). But a $*$ point of $N[Y \setminus V]$ is a point of NY which — because (21) is cartesian — is not a point of NV , a contradiction to the fact that $NY \subseteq NZ \subseteq NV$ by assumption.

So without loss of generality (by replacing X with V), we can assume that Z is a *closed* subscheme of X . Factoring $Y \hookrightarrow X$ as $Y \xrightarrow{i_Y} U \xrightarrow{j_Y} X$ with a closed immersion i_Y and an open immersion j_Y and replacing X with U and Z with $Z \cap U$, we can furthermore assume that Y is also a closed subscheme of X .

Finally, since the question is local on X , we can assume that $X = \operatorname{Spec}(A)$ is affine and that Y and Z are given by ideals I and J of A . By assumption, we have $J \cdot NA \subseteq I \cdot NA$, and using 6.3, we conclude

$$J = A \cap [J \cdot NA] \subseteq A \cap [I \cdot NA] = I.$$

q.e.d.

6.14. Remark. Let \mathcal{C} be a category with fibred products and a terminal object T , let X and Y be two objects of \mathcal{C} , and let $f, g : X \rightarrow Y$ be two morphisms. Then the *equalizer*

$$\operatorname{Eq}(f, g) \xrightarrow{\operatorname{eq}(f, g)} X \xrightleftharpoons[g]{f} Y$$

of f and g exists — it is given by the cartesian diagram

$$(22) \quad \begin{array}{ccc} \operatorname{Eq}(f, g) & \xrightarrow{\operatorname{eq}(f, g)} & X \\ \downarrow & \square & \downarrow (f, g) \\ Y & \xrightarrow{\langle \mathbb{1}_Y, \mathbb{1}_Y \rangle} & Y \times_T Y \end{array}$$

6.15. Lemma. Let S be a scheme, let X and Y be two S -schemes, and let $f, g : X \rightarrow Y$ be two S -morphisms. Then the equalizer

$$\operatorname{Eq}(f, g) \xrightarrow{\operatorname{eq}(f, g)} X \xrightleftharpoons[g]{f} Y$$

of f and g exists in the category of S -schemes and is an immersion.

Proof. The category of S -schemes has fibred products and the terminal object S , so the equalizer of f and g exists by 6.14. It is an immersion by the construction given in (22), because $Y \xrightarrow{\langle \mathbb{1}_Y, \mathbb{1}_Y \rangle} Y \times_S Y$ is an immersion. **q.e.d.**

6.16. Corollary. The functor $N : \operatorname{Sch}_k^{\operatorname{fp}} \rightarrow * \operatorname{Sch}_k^{\operatorname{fp}}$ is faithful.

Proof. Let X and Y be k -schemes of finite type, and let $f, g : X \rightarrow Y$ be k -morphisms with $Nf = Ng$. By 6.16, f and g are equal if and only if $\operatorname{Eq}(f, g)$ equals X as subschemes of X . By assumption, $\operatorname{Eq}(Nf, Ng)$ is the $*$ subscheme NX of NX , and $\operatorname{Eq}(Nf, Ng) = N\operatorname{Eq}(f, g)$ by 4.1(i), so the claim follows from 6.13. **q.e.d.**

Let S be a noetherian scheme, let X/S be projective with very ample sheaf $\mathcal{O}(1)$, let \mathcal{F} be a coherent sheaf on X , and let $P \in \mathbb{Q}[t]$ be a rational polynomial. Then we have the *Quot-scheme* $\text{Quot}_{\mathcal{F}/X/S}^P$, projective over S , which represents the contravariant functor $T \mapsto \text{Quot}^P(\mathcal{F}_{X \times_S T}/X_T/T)$ that maps a locally noetherian S -scheme T to the set of those quotients $\mathcal{F}_{X \times_S T} \twoheadrightarrow \mathcal{G}$ with \mathcal{G} flat over T and Hilbert polynomial P in every fibre $t \in T$ (compare [FGA, 221.3]).

By transfer, for a \ast noetherian \ast scheme S , a \ast projective S - \ast scheme X with \ast very ample \ast sheaf $\mathcal{O}(1)$, a \ast coherent \ast sheaf \mathcal{F} on X and a \ast polynomial $P \in \ast\mathbb{Q}^\ast[t]$, we have a canonical \ast projective S - \ast scheme $\ast\text{Quot}_{\mathcal{F}/X/S}^P$ which represents the enlarged functor $T \mapsto \ast\text{Quot}^P(\mathcal{F}_{X \times_S T}/X_T/T)$ on \ast locally noetherian S - \ast schemes.

In the special case $\mathcal{F} = \mathcal{O}_X$, the Quot-scheme $\text{Quot}_{\mathcal{O}_X/X/S}^P$ is called the *Hilbert scheme* and denoted by $\text{Hilb}_{X/S}^P$ (its T -valued points correspond to closed subschemes of X_T which are flat over T and have Hilbert polynomial P in every fibre). — Similarly, we call $\ast\text{Hilb}_{X/S}^P := \ast\text{Quot}_{\mathcal{O}_X/X/S}^P$ the \ast Hilbert scheme.

In the following proposition, we want to show that the formation of Quot-schemes and Hilbert schemes is compatible with the functor N :

6.17. Proposition. Let X be a projective k -scheme with very ample sheaf $\mathcal{O}(1)$, let \mathcal{F} be a coherent sheaf on X , and let $P \in \mathbb{Q}[t]$ be a rational polynomial.

- (i) We have $N \text{Quot}_{\mathcal{F}/X/k}^P = \ast\text{Quot}_{N\mathcal{F}/N X/k}^P$ and in particular $N \text{Hilb}_{X/k}^P = \ast\text{Hilb}_{N X/k}^P$, where P is considered as a \ast polynomial via $\mathbb{Q}[t] \hookrightarrow \ast\mathbb{Q}^\ast[t]$.
- (ii) Let T be a k -scheme of finite type, and let $f : T \rightarrow \text{Quot}_{\mathcal{F}/X/k}^P$ be a T -valued point, corresponding to a quotient $\varphi : \mathcal{F}_{X \times_k T} \twoheadrightarrow \mathcal{G}$. Then $[Nf]$, which is a $[NT]$ -valued point of $\ast\text{Quot}_{N\mathcal{F}/N X/k}^P$ by (i), corresponds to the quotient $[N\mathcal{F}]_{[NX] \times_k [NT]} \twoheadrightarrow N\mathcal{G}$.

In particular, if $g : T \rightarrow \text{Hilb}_{X/k}^P$ corresponds to the subscheme $Z \subseteq X \times_k T$, then $[Ng]$ corresponds to the \ast subscheme $[NZ] \subseteq [NX] \times_k [NT]$.

Proof. By [EGA IV₃, 8.5.2, 8.8.2, 8.10.5], there exist a finitely generated subring A_0 of k , a projective A_0 -scheme X_0 with $X = X_0 \times_{A_0} k$ and a coherent sheaf \mathcal{F}_0 on X_0 with $[A_0 \hookrightarrow k]^\ast \mathcal{F}_0 = \mathcal{F}$.

Then $\text{Quot}_{\mathcal{F}/X/k}^P = \text{Quot}_{\mathcal{F}_0/X_0/A_0}^P \times_{A_0} k$, and putting $\alpha := \tau_{\mathbb{Z}, A_0, k}^{-1}[A_0 \hookrightarrow k] : \ast A_0 \rightarrow k$, we get

$$\begin{aligned} N \text{Quot}_{\mathcal{F}/X/k}^P &\stackrel{3.4}{=} \alpha^\ast \left(\ast [\text{Quot}_{\mathcal{F}_0/X_0/A_0}^P] \right) = \alpha^\ast \left[\ast \text{Quot}_{\ast \mathcal{F}_0/\ast X_0/\ast A_0}^P \right] \\ &= \ast \text{Quot}_{\alpha^\ast[\ast \mathcal{F}_0]/\alpha^\ast[\ast X_0]/\alpha^\ast[\ast A_0]}^P \stackrel{3.4.5.2}{=} \ast \text{Quot}_{N\mathcal{F}/N X/k}^P, \end{aligned}$$

which settles (i).

By [EGA IV₃, 8.8.2], after a possible change of A_0 , X_0 and \mathcal{F}_0 , we find an A_0 -scheme T_0 of finite type with $T = T_0 \times_{A_0} k$ and an A_0 -morphism $f_0 : T_0 \rightarrow Q := \text{Quot}_{\mathcal{F}_0/X_0/A_0}^P$ with $[A_0 \hookrightarrow k]^\ast f_0 = f$. Let $\mathcal{F}_{X_0 \times_{A_0} Q} \twoheadrightarrow \mathcal{G}^{\text{univ}}$ be the universal quotient. Then f_0 corresponds to the quotient $[\mathcal{F}_0]_{X_0 \times_{A_0} T_0} \twoheadrightarrow [\mathbb{1}_{X_0} \times f_0]^\ast \mathcal{G}^{\text{univ}} =: \mathcal{G}_0$, and f corresponds to the quotient $\varphi : \mathcal{F}_{X \times_k T} \twoheadrightarrow [A_0 \hookrightarrow k]^\ast \mathcal{G}_0 = \mathcal{G}$. So

$$N \mathcal{G} \stackrel{5.2}{=} \alpha^\ast (\ast \mathcal{G}_0) = \alpha^\ast \left(\ast [\mathbb{1}_{X_0} \times f_0]^\ast \mathcal{G}^{\text{univ}} \right) \stackrel{3.4}{=} [\mathbb{1}_{NX} \times [Nf]]^\ast (\alpha^\ast (\ast \mathcal{G}^{\text{univ}})),$$

and since $[N\mathcal{F}]_{[NX] \times_k \ast \text{Quot}_{N\mathcal{F}/N X/k}^P} \twoheadrightarrow \alpha^\ast (\ast \mathcal{G}^{\text{univ}})$ obviously is the universal quotient, this proves (ii).

q.e.d.

6.18. Corollary. Let X be a projective k -scheme with very ample sheaf $\mathcal{O}(1)$, and let \mathcal{F} be a coherent \mathcal{O}_X -module.

Then the Hilbert polynomial of \mathcal{F} (with respect to $\mathcal{O}(1)$) coincides with the \ast Hilbert polynomial of $N\mathcal{F}$ (with respect to $N\mathcal{O}(1)$) in $\ast\mathbb{Q}^\ast[t]$.

Proof. Denote the Hilbert polynomial of \mathcal{F} by $P_{\mathcal{F}} \in \mathbb{Q}[t] \subset \ast\mathbb{Q}^\ast[t]$. If \mathcal{F} corresponds to the k -valued point f of $\text{Quot}_{\mathcal{F}/X/k}^{P_{\mathcal{F}}}$, then $N\mathcal{F}$ corresponds to the k -valued point $[Nf]$ of $\ast\text{Quot}_{N\mathcal{F}/N X/k}^{P_{\mathcal{F}}}$ according to 6.17(ii). But by its very definition, $\ast\text{Quot}_{N\mathcal{F}/N X/k}^{P_{\mathcal{F}}}$ parameterizes sheaves with \ast Hilbert polynomial $P_{\mathcal{F}} \in \ast\mathbb{Q}^\ast[t]$, so we are done. **q.e.d.**

6.19. Theorem. Let X be a projective k -scheme with very ample sheaf $\mathcal{O}(1)$, and let \mathcal{G} be a $*$ coherent $*$ sheaf on NX . Then the following two statements are equivalent:

- (i) There is a coherent sheaf \mathcal{H} on X with $N\mathcal{H} \cong \mathcal{G}$.
- (ii) There is a coherent sheaf \mathcal{F} on X , such that \mathcal{G} is a quotient of $N\mathcal{F}$, and the $*$ Hilbert polynomial of \mathcal{G} (with respect to $N\mathcal{O}(1)$) lies in $\mathbb{Q}[t] \subset *Q^*[t]$.

Proof. The implication “(i) \Rightarrow (ii)” is easy: We can simply put $\mathcal{F} := \mathcal{H}$, and by 6.18, the $*$ Hilbert polynomial of $\mathcal{G} \cong N\mathcal{H}$ equals the Hilbert polynomial of \mathcal{H} and consequently lies in $\mathbb{Q}[t]$.

For “(ii) \Rightarrow (i)”, let $P \in \mathbb{Q}[t] \subset *Q^*[t]$ be the $*$ Hilbert polynomial of \mathcal{G} . Then $N\mathcal{F} \twoheadrightarrow \mathcal{G}$ corresponds to a k -valued point g of $*\text{Quot}_{N\mathcal{F}/NX/k}^P$. Since

$$\left[\text{Quot}_{\mathcal{F}/X/k}^P \right](k) \xrightarrow{N} \left[N \text{Quot}_{\mathcal{F}/X/k}^P \right](k) \stackrel{6.17(i)}{=} \left[* \text{Quot}_{N\mathcal{F}/NX/k}^P \right](k)$$

is bijective by 4.13, there exists a k -valued point h of $\text{Quot}_{\mathcal{F}/X/k}^P$ with $g = Nh$. If $\mathcal{F} \twoheadrightarrow \mathcal{H}$ is the quotient given by h , then $N\mathcal{H} \cong \mathcal{G}$ by 6.17(ii). **q.e.d.**

6.20. Corollary. Let X be a projective k -scheme with very ample sheaf $\mathcal{O}(1)$, and let Z be a $*$ closed $*$ subscheme of NX . Then the following two statements are equivalent:

- (i) There is a closed subscheme W of X with $NW = Z$.
- (ii) The $*$ Hilbert polynomial of Z (with respect to $N\mathcal{O}(1)$) lies in $\mathbb{Q}[t] \subset *Q^*[t]$.

Proof. This follows immediately from 6.19, applied to the special case $\mathcal{G} := \mathcal{O}_Z$ and $\mathcal{F} := \mathcal{O}_X$. **q.e.d.**

6.21. Corollary. Let X be a projective k -scheme with very ample sheaf $\mathcal{O}(1)$, and let Z be a $*$ closed $*$ integral $*$ subscheme (i.e. a $*$ prime cycle) of NX that has *finite* $*$ degree (with respect to $[Z \hookrightarrow NX]^* \mathcal{O}(1)$). Then there exists an integral subscheme (i.e. a prime cycle) W of X with $NW = Z$.

Proof. As Z is a subscheme of NX , we have $*\dim Z \leq *\dim NX \stackrel{6.4(ii)}{=} \dim X$, so Z is a $*$ projective $*$ integral $*$ scheme of finite $*$ degree and of *finite* $*$ dimension. Then transfer of [SGA 6, XIII.6.11(i)] shows that the $*$ Hilbert polynomial of Z has finite coefficients and consequently lies in $\mathbb{Q}[t] \subset *Q^*[t]$, and the corollary follows from 6.20. **q.e.d.**

6.22. Corollary. Let $n \in \mathbb{N}_+$, and let Z be a $*$ integral $*$ closed $*$ subscheme of $*\mathbb{P}_k^n$ of finite $*$ degree. Then there is an integral closed subscheme W of \mathbb{P}_k^n with $NW = Z$.

Proof. This follows immediately from 6.21 for $X := \mathbb{P}_k^n$ and $\mathcal{O}(1) := N\mathcal{O}_{\mathbb{P}_k^n}(1)$. **q.e.d.**

Let S be a scheme, and let $f : X \rightarrow Y$ be an S -morphism. Then the *graph of f* is the S -morphism $\Gamma_f : X \xrightarrow{\langle \mathbb{1}_X, f \rangle} X \times_S Y$. It is easy to see that the diagram

$$\begin{array}{ccc} X & \xrightarrow{\Gamma_f} & X \times_S Y \\ f \downarrow & \square & \downarrow f \times \mathbb{1}_Y \\ Y & \xrightarrow{\langle \mathbb{1}_Y, \mathbb{1}_Y \rangle} & Y \times_S Y \end{array}$$

is cartesian, which shows that Γ_f is an immersion (and can hence be considered as a subscheme of $X \times_S Y$, isomorphic to X), which is closed if Y/S is separated.

Now let S be noetherian, let X and Y be projective S -schemes with X/S flat, let $\mathcal{O}(1)$ be a very ample sheaf on $X \times_S Y$, and let $P \in \mathbb{Q}[t]$ be a polynomial. Consider the functor $T \mapsto \text{Hom}_S^P(X, Y)(T)$ that

maps an S -scheme T to the set of those T -morphisms $f : X \times_S T \rightarrow Y \times_S T$ whose graph $\Gamma_f \hookrightarrow X \times_S Y$, a closed subscheme since Y/S is separated, has Hilbert polynomial P with respect to $\mathcal{O}(1)$.

It is well known (compare [Kol96, I.1.10]) that this functor is represented by an open subscheme $\mathrm{Hom}_S^P(X, Y)$ of $\mathrm{Hilb}_{X \times_S Y/S}^P$, where $\mathrm{Hom}_S^P(X, Y) \hookrightarrow \mathrm{Hilb}_{X \times_S Y/S}^P$ is given by sending a morphism to its graph. Similar to the case of Quot- and Hilbert schemes, the formation of $\mathrm{Hom}_S^P(X, Y)$ is compatible with the functor N in the following sense:

6.23. Proposition. Let X and Y be projective k -schemes, let $\mathcal{O}(1)$ be a very ample sheaf on $X \times_k Y$, and let $P \in \mathbb{Q}[t]$ be a rational polynomial.

- (i) We have $N \mathrm{Hom}_k^P(X, Y) = {}^* \mathrm{Hom}_k^P(NX, NY)$, where P is considered as a $*$ polynomial via $\mathbb{Q}[t] \hookrightarrow {}^* \mathbb{Q}^*[t]$.
- (ii) Let T be a k -scheme of finite type, and let $f : T \rightarrow \mathrm{Hom}_k^P(X, Y)$ be a T -valued point, corresponding to a T -morphism $g : X \times_k T \rightarrow Y \times_k T$. Then $[Nf]$, which is a $[NT]$ -valued point of ${}^* \mathrm{Hom}_k^P(NX, NY)$ by (i), corresponds to the morphism $[Ng] : NX^* \times_k NT \rightarrow NY^* \times_k NT$.

Proof. This is completely analogous to the proof of 6.17. **q.e.d.**

6.24. Theorem. Let X and Y be projective k -schemes, let $\mathcal{O}(1)$ be a very ample sheaf on $X \times_k Y$, and let $g : NX \rightarrow NY$ be a morphism of k - $*$ schemes. Then the following two statements are equivalent:

- (i) There is a k -morphism $f : X \rightarrow Y$ with $Nf = g$.
- (ii) The $*$ Hilbert polynomial of the $*$ graph of g (with respect to $N\mathcal{O}(1)$) lies in $\mathbb{Q}[t] \subset {}^* \mathbb{Q}^*[t]$.

Proof. This follows from 6.23 in the same way as 6.19 follows from 6.17. **q.e.d.**

6.25. Corollary. Let X and Y be projective k -schemes with X integral, let $\mathcal{O}(1)$ be a very ample sheaf on $X \times_k Y$, and let $g : NX \rightarrow NY$ be a morphism of k - $*$ schemes whose $*$ graph has *finite* degree (with respect to $N\mathcal{O}(1)$). Then there exists a k -morphism $f : X \rightarrow Y$ with $Nf = g$.

Proof. By transfer, the $*$ graph ${}^* \Gamma_g$ of g is isomorphic to NX and hence $*$ integral. Then by 6.21, there is a closed subscheme Γ of $X \times_k Y$ with $N\Gamma = {}^* \Gamma_g$, and it follows from 6.20 that the $*$ Hilbert polynomial of ${}^* \Gamma_g$ lies in $\mathbb{Q}[t]$. Then the corollary follows from 6.24. **q.e.d.**

6.26. Corollary. The restriction of $N : \mathcal{Sch}_k^{\mathrm{fp}} \rightarrow {}^* \mathcal{Sch}_k^{\mathrm{fp}}$ to the full subcategory of projective k -schemes reflects isomorphisms.

Proof. Let $f : X \rightarrow Y$ be a morphism of projective k -schemes such that Nf is an isomorphism with inverse $\tilde{g} : NY \rightarrow NX$. Choose a very ample sheaf $\mathcal{O}(1)$ on $X \times_k Y$. If $\tau : Y \times_k X \xrightarrow{\sim} X \times_k Y$ denotes the transposition, $\tau^* \mathcal{O}(1)$ is a very ample sheaf on $Y \times_k X$. Let $P \in \mathbb{Q}[t]$ be the Hilbert polynomial of Γ_f with respect to $\mathcal{O}(1)$, which by 6.18 is also the $*$ Hilbert polynomial of ${}^* \Gamma_{Nf}$, the $*$ graph of Nf , with respect to $N\mathcal{O}(1)$. It follows from transfer that the transpose $[N\tau]^* [{}^* \Gamma_{Nf}]$ is the $*$ graph of \tilde{g} and that its $*$ Hilbert polynomial with respect to $N[\tau^* \mathcal{O}(1)]$ equals P . Thus by 6.24, there is a k -morphism $g : Y \rightarrow X$ with $Ng = \tilde{g}$. Now

$$N[f \circ g] = [Nf] \circ [Ng] = [Nf] \circ \tilde{g} = \mathbb{1}_{NY} = N\mathbb{1}_Y$$

and

$$N[g \circ f] = [Ng] \circ [Nf] = \tilde{g} \circ [Nf] = \mathbb{1}_{NX} = N\mathbb{1}_X,$$

so $f \circ g = \mathbb{1}_Y$ and $g \circ f = \mathbb{1}_X$ (because $N : \mathcal{Sch}_k^{\mathrm{fp}} \rightarrow {}^* \mathcal{Sch}_k^{\mathrm{fp}}$ is faithful by 6.16), and we see that f is indeed an isomorphism (with inverse g). **q.e.d.**

6.27. Lemma. Let $\varphi : B \hookrightarrow C$ be a finite, injective morphism of integral k -algebras of finite type. Then $N\varphi : NB \rightarrow NC$ is an injective, finite morphism of integral k -algebras, and

$$[\text{Quot}(NC) : \text{Quot}(NB)] = [\text{Quot}(C) : \text{Quot}(B)] \in \mathbb{N}_+.$$

Proof. The (internal) k -algebras NB and NC are integral by 6.4(iii), and $N\varphi$ is injective and finite, because

$$\begin{array}{ccc} B & \xrightarrow{\varphi} & C \\ \sigma_B \downarrow & & \downarrow \sigma_C \\ NB & \xrightarrow{N\varphi} & NC \end{array}$$

is cocartesian by 4.16 and because σ_B is faithfully flat by 6.3. Since $\text{Quot}(C) = C \otimes_B \text{Quot}(B)$ and

$$\begin{aligned} \text{Quot}(NC) &= [NC] \otimes_{NB} \text{Quot}(NB) \stackrel{4.16}{=} C \otimes_B \text{Quot}(NB) \\ &= [C \otimes_B \text{Quot}(B)] \otimes_{\text{Quot}(B)} \text{Quot}(NB) = \text{Quot}(C) \otimes_{\text{Quot}(B)} \text{Quot}(NB). \end{aligned}$$

Using this, we get

$$\begin{aligned} [\text{Quot}(NC) : \text{Quot}(NB)] &= \dim_{\text{Quot}(NB)} \text{Quot}(NC) \\ &= \dim_{\text{Quot}(NB)} [\text{Quot}(C) \otimes_{\text{Quot}(B)} \text{Quot}(NB)] = \dim_{\text{Quot}(B)} \text{Quot}(C) = [\text{Quot}(C) : \text{Quot}(B)], \end{aligned}$$

and this degree is of course finite, because φ is finite. **q.e.d.**

6.28. Proposition. Let $f : X \rightarrow Y$ be a morphism of integral k -schemes of finite type. Then f is birational if and only if $Nf : NX \rightarrow NY$ is $*$ birational.

Proof. Assume first that f is birational. Then by definition, there is a commutative diagram

$$\begin{array}{ccc} & U & \\ j_1 \swarrow & & \searrow j_2 \\ X & \xrightarrow{f} & Y \end{array}$$

of k -morphisms with open immersions j_1 and j_2 . So

$$\begin{array}{ccc} & NU & \\ Nj_1 \swarrow & & \searrow Nj_2 \\ NX & \xrightarrow{Nf} & NY \end{array}$$

is a commutative diagram of k - $*$ schemes, where Nj_1 and Nj_2 are $*$ open immersions by 4.4, which shows that Nf is $*$ birational.

For the other implication, assume now that Nf is $*$ birational. Then NX and NY have the same $*$ dimension, and 6.4(ii) implies that $\dim X = \dim Y$. Let us first show that f is dominant: If it were not, there would be a non-empty open subscheme U of Y and a cartesian diagram

$$\begin{array}{ccc} \emptyset & \longrightarrow & U \\ \downarrow & \square & \downarrow \\ X & \xrightarrow{f} & Y. \end{array}$$

But then by 4.1(i), 4.4 and 6.4(i), NU would be a non-empty $*$ open $*$ subscheme of NY disjoint from $[Nf](NX)$; this means that Nf would not be $*$ dominant and consequently could not be $*$ birational — a contradiction. So f is indeed dominant, and if we denote the generic points of X and Y by ξ respectively η , then ξ is contained in the generic fibre X_η . Since we saw above that $\dim X = \dim Y$, we must have $X_\eta = \{\xi\}$ by [EGA IV₂, 4.1.2(i)].

In particular, X_η/η is of finite type and discrete, so by [EGA I, 6.4.4] it is finite. Then by [EGA IV₃, p. 6 and 8.10.5(x)], there is an affine, open, dense subset $V = \text{Spec}(B) \subseteq Y$, such that $f|_U : U \rightarrow V$

(with $U := f^{-1}(V)$) is finite. Then $U = \operatorname{Spec}(C)$ is affine, and $f^* : B \rightarrow C$ is a finite, injective morphism of integral k -algebras of finite type. By hypothesis we have $\operatorname{Quot}(NB) \xrightarrow{\sim} \operatorname{Quot}(NC)$, so 6.27 implies $k(Y) = \operatorname{Quot}(B) \xrightarrow{\sim} \operatorname{Quot}(C) = k(X)$, which means that f induces an isomorphism of the function fields of X and Y and is therefore birational. **q.e.d.**

7. THE COHERENCE THEOREM

For any scheme X , sheaf of \mathcal{O}_X -modules \mathcal{F} and natural number $i \in \mathbb{N}_0$, we can consider the Zariski cohomology group $H^i(X, \mathcal{F})$. If X is an A -scheme for a ring A , then $H^i(X, \mathcal{F})$ canonically carries the structure of an A -module.

If $f : X \rightarrow Y$ is a proper morphism of schemes and if \mathcal{F} is a coherent \mathcal{O}_X -module, then we have the higher direct image $R^i f_* \mathcal{F}$, a coherent \mathcal{O}_Y -module by [EGA III₁, 3.2.1].

By transfer, if X is a $*$ scheme, \mathcal{F} a $*$ finitely presented \mathcal{O}_X -module and $i \in \mathbb{N}_0$ a $*$ natural number, we get the *$*$ Zariski cohomology* $H^i(X, \mathcal{F})$ which is an internal A -module if X is an A - $*$ scheme for a $*$ ring A .

Similarly, if $f : X \rightarrow Y$ is a $*$ proper morphism of $*$ schemes and if \mathcal{F} is a $*$ coherent \mathcal{O}_X -module, we have the *$*$ higher direct image* $R^i f_* \mathcal{F}$, a $*$ coherent \mathcal{O}_Y -module.

7.1. Lemma. Let A be a $*$ noetherian $*$ ring, and let $f : X \rightarrow Y$ be a morphism of $*$ schemes over A . Then the left exact functor $f_* : *QCoh_X \rightarrow *Mod_Y$ factorizes over $*QCoh_Y$ and admits a right derived functor $R(f)_* : D^+(*QCoh_X) \rightarrow D^+(*QCoh_Y)$.

Furthermore, the class of flasque[†] $*$ quasi-coherent sheaves of \mathcal{O}_X - $*$ modules is adapted to f_* .

Proof. Let B be a noetherian ring in \mathcal{R} , and let $g : Z \rightarrow W$ be a morphism of finitely presented B -schemes. Then Z and g are quasi-separated (by [EGA IV₁, 1.2.8]) and quasi-compact. It follows from [TT90, B.3] that $QCoh_Z^u$ has enough injective objects and from [TT90, B.6] that $R^i g_* \mathcal{F}$ is quasi-coherent for all quasi-coherent \mathcal{O}_Z -modules \mathcal{F} and all $i \in \mathbb{N}_0$.

Furthermore, by [TT90, B.4], an injective object in $QCoh_Z^u$ is also an injective (and hence flasque) object of Mod_Z^u , so that the class of flasque quasi-coherent \mathcal{O}_Z -modules is adapted to g_* .

Since all this is true for arbitrary B , Z , W and g , the transferred statements are also true, and the lemma follows. **q.e.d.**

7.2. Lemma. Let A be a $*$ noetherian $*$ ring, and let $f : X \rightarrow Y$ be a morphism of finitely presented A -schemes. Then the following diagram of exact functors between derived categories commutes (up to canonical isomorphism):

$$\begin{array}{ccc} D^+(*QCoh_{NX}) & \xrightarrow{R[Nf]_*} & D^+(*QCoh_{NY}) \\ \downarrow S & & \downarrow S \\ D^+(Mod_X) & \xrightarrow{Rf_*} & D^+(Mod_Y). \end{array}$$

Proof. First of all, note that $R[Nf]_*$ exists by 7.1 and that $S : *QCoh_{NX} \rightarrow Mod_X$ and $S : *QCoh_{NY} \rightarrow Mod_Y$ are exact by 5.8.

The composition $S \circ R[Nf]_*$ is canonically isomorphic to $R[S \circ [Nf]_*]$, because S is exact. The composition $Rf_* \circ S$ is canonically isomorphic to $R[f_* \circ S]$, because S is exact and obviously maps flasque $*$ sheaves to flasque sheaves, which are adapted to f_* .

It follows immediately from the definition of S , f_* and $[Nf]_*$ that $S \circ [Nf]_* = f_* \circ S$, so we have

$$S \circ R[Nf]_* \cong R[S \circ [Nf]_*] = R[f_* \circ S] \cong Rf_* \circ S.$$

[†]Note that being flasque is obviously first-order and hence is the same as being $*$ flasque.

q.e.d.

Let k be a $*$ field, and let $f : X \longrightarrow Y$ be a *proper* morphism of k -schemes of finite type.

7.3. Lemma. We have a commutative diagram of exact functors

$$\begin{array}{ccc}
 D^b(*\mathcal{Coh}_{NX}) & \xrightarrow{R[Nf]_*} & D^b(*\mathcal{Coh}_{NY}) \\
 \downarrow \iota & & \downarrow \iota \\
 D^b(*Q\mathcal{Coh}_{NX}) & \xrightarrow{R[Nf]_*} & D^b(*Q\mathcal{Coh}_{NY}) \\
 \downarrow S & & \downarrow S \\
 D^b(\mathcal{Mod}_X) & \xrightarrow{Rf_*} & D^b(\mathcal{Mod}_Y) \\
 \downarrow \iota & & \downarrow \iota \\
 D^b(\mathcal{Coh}_X) & \xrightarrow{Rf_*} & D^b(\mathcal{Coh}_Y)
 \end{array}$$

Proof. Since X is finitely presented over a field, it is finite-dimensional, which implies that $f_* : Q\mathcal{Coh}_X \rightarrow Q\mathcal{Coh}_Y$ has finite cohomological dimension and hence induces $Rf_* : D^b(Q\mathcal{Coh}_X) \rightarrow D^b(Q\mathcal{Coh}_Y)$.

By 6.4(ii) and transfer, $[Nf]_* : *Q\mathcal{Coh}_{NX} \rightarrow *Q\mathcal{Coh}_{NY}$ has the same finite cohomological dimension and induces $R[Nf]_* : D^b(*Q\mathcal{Coh}_{NX}) \rightarrow D^b(*Q\mathcal{Coh}_{NY})$. So the middle square is well-defined, and it commutes by 7.2.

The bottom square is well-defined and commutes by [Har66, II.2.2] and [SGA 6, II.2.2.2], the top square is well-defined and commutes by transfer of [Har66, II.2.2] and [SGA 6, II.2.2.1]. **q.e.d.**

7.4. Proposition. There is a canonical morphism of exact functors

$$(23) \quad \begin{array}{ccc}
 & \xrightarrow{\iota \circ Rf_*} & \\
 D^b(\mathcal{Coh}_X) & \Downarrow & D^b(\mathcal{Mod}_Y) \\
 & \xleftarrow{S \circ R[Nf]_* \circ N} &
 \end{array}$$

which induces a canonical morphism of δ -functors

$$(24) \quad \begin{array}{ccc}
 & \xrightarrow{(N \circ R^n f_*)_{n \in \mathbb{N}_0}} & \\
 \mathcal{Coh}_X & \Downarrow & *\mathcal{Coh}_{NY} \\
 & \xleftarrow{(R^n [Nf]_* \circ N)_{n \in \mathbb{N}_0}} &
 \end{array}$$

Proof. Morphism (23) is given by the following diagram in the 2-category of triangulated categories

$$\begin{array}{ccccc}
 & & D^b(\mathcal{Coh}_Y) & & \\
 & \nearrow Rf_* & \downarrow & \searrow \iota & \\
 D^b(\mathcal{Coh}_X) & \xrightarrow{\iota} & D^b(\mathcal{Mod}_X) & & D^b(\mathcal{Mod}_Y) \\
 \downarrow N & \downarrow & \downarrow & \downarrow Rf_* & \downarrow S \\
 & D^b(*\mathcal{Coh}_{NX}) & & D^b(*\mathcal{Coh}_{NY}) & \\
 & \nearrow Rf_* & & \nearrow S &
 \end{array}$$

where the three 2-morphisms are given by (9) and 7.3 (note that $N : \mathcal{Coh}_X \rightarrow {}^*\mathcal{Coh}_{NX}$ is exact by 6.4(iv)).

Applying (23) to objects concentrated in degree zero (i.e. objects coming from \mathcal{Coh}_X) and taking cohomology gives us a morphism of δ -functors

$$\begin{array}{ccc} & \xrightarrow{(R^n f_*)_{n \in \mathbb{N}_0}} & \\ \mathcal{Coh}_X & \Downarrow \varphi & \mathcal{Mod}_Y \\ & \xrightarrow{(S \circ R^n[Nf]_* \circ N)_{n \in \mathbb{N}_0}} & \end{array}$$

Using 5.12, we then get the morphism from (24) for a coherent \mathcal{O}_X -module \mathcal{F} and an $n \in \mathbb{N}_0$ by

$$N R^n f_* \mathcal{F} \xrightarrow{\tau_{Y, R^n f_* \mathcal{F}, R^n[Nf]_* N \mathcal{F}}^{-1}(\varphi)} R^n[Nf]_* N \mathcal{F}.$$

That this is indeed a morphism of δ -functors follows immediately from the exactness of N , from 5.13 and from the fact that φ is a morphism of δ -functors. **q.e.d.**

7.5. Theorem. The canonical morphism of functors (24) is an isomorphism. In particular, $N R^n f_* \mathcal{F}$ is canonically isomorphic to $R^n[Nf]_* N \mathcal{F}$ for all coherent \mathcal{O}_X -modules \mathcal{F} and all $n \in \mathbb{Z}$.

Proof. Because the statement is local in Y , we can assume without loss of generality that $Y = \text{Spec}(B)$ is affine for a finitely presented A -algebra B . We split the proof in several cases:

First consider the case where $f : X = \mathbb{P}_Y^d \rightarrow Y$ is the structural morphism of *projective d -space* over Y . By [EGA III₁, 2.1.15, 2.1.16], for any $m, n \in \mathbb{Z}$, we have canonical isomorphisms

$$R^n f_* \mathcal{O}_X(m) = \begin{cases} \mathcal{O}_Y[T_0, \dots, T_d]_m & \text{if } n = 0, \\ \mathcal{O}_Y[T_0, \dots, T_d]_{-d-1-m}^\vee & \text{if } n = d, \\ 0 & \text{otherwise,} \end{cases}$$

where $\mathcal{O}_Y[T_0, \dots, T_d]$ denotes the graded free symmetric algebra over \mathcal{O}_Y with generators T_0, \dots, T_d (so that its part of degree m is just the free \mathcal{O}_Y -module with basis the homogenous monomials of degree m in the T_i). By 3.7, 5.18(i) and transfer, we have

$$R^n[Nf]_*[N \mathcal{O}_X(m)] = \begin{cases} \mathcal{O}_{NY}^*[T_0, \dots, T_d]_m & \text{if } n = 0, \\ \mathcal{O}_{NY}^*[T_0, \dots, T_d]_{-d-1-m}^\vee & \text{if } n = d, \\ 0 & \text{otherwise.} \end{cases}$$

Since a $*$ monomial of degree m is the same as a monomial of degree m , and since N respects duals by 5.17, we see that $N R^n f_* \mathcal{O}_X(m) = R^n[Nf]_*[N \mathcal{O}_X(m)]$ for all m and n . By additivity, the theorem is hence true for our special choice of f and for all \mathcal{F} of the form $\mathcal{O}_X(m)^l$ for $l \in \mathbb{N}_0$ and $m \in \mathbb{Z}$.

As a next step, we prove the theorem for all coherent sheaves on \mathbb{P}_Y^d by decreasing induction on n (this part closely resembles Hartshorne's proof of the "Theorem on Formal Functions" in [Har93]): Since $R^n f_*$ and $R^n[Nf]_*$ both vanish for $n > d$, the theorem holds trivially in those cases. For the inductive step, assume that the theorem holds for all $n' > n \in \mathbb{N}_0$, and let \mathcal{F} be an arbitrary coherent sheaf on X . By [EGA III₁, 2.2.2(iv)], there exists an epimorphism $\mathcal{G} := \mathcal{O}_X(m)^l \twoheadrightarrow \mathcal{F}$ for suitable $l \in \mathbb{N}_0$ and $m \in \mathbb{Z}$, so that we have a short exact sequence

$$0 \longrightarrow \mathcal{H} \longrightarrow \mathcal{G} \longrightarrow \mathcal{F} \longrightarrow 0$$

of coherent \mathcal{O}_X -modules. By 7.4, we get an induced commutative diagram of $*$ coherent \mathcal{O}_{NX} -modules with exact rows as follows:

$$\begin{array}{ccccccccc} N R^n f_* \mathcal{H} & \longrightarrow & N R^n f_* \mathcal{G} & \longrightarrow & N R^n f_* \mathcal{F} & \longrightarrow & N R^{n+1} f_* \mathcal{H} & \longrightarrow & N R^{n+1} f_* \mathcal{G} \\ \alpha \downarrow & & \beta \downarrow \wr & & \gamma \downarrow \cdots & & \delta \downarrow \wr & & \varepsilon \downarrow \wr \\ R^n[Nf]_*[N \mathcal{H}] & \longrightarrow & R^n[Nf]_*[N \mathcal{G}] & \longrightarrow & R^n[Nf]_*[N \mathcal{F}] & \longrightarrow & R^{n+1}[Nf]_*[N \mathcal{H}] & \longrightarrow & R^{n+1}[Nf]_*[N \mathcal{G}] \end{array}$$

By the first part of the proof, β and ε are isomorphisms, and by our inductive hypothesis, δ is an isomorphism. Then by the five lemma, since β and δ are epimorphisms and ε is a monomorphism, γ is an epimorphism.

Since \mathcal{F} was chosen arbitrarily, this conclusion also applies to \mathcal{H} , i.e. α is also an epimorphism. But then we can apply the five lemma again, using that α is an epimorphism and that β and δ are monomorphisms, to conclude that γ is a monomorphism and hence an isomorphism as desired.

Having settled the theorem for projective space, we now consider the **second case** where $f : X \hookrightarrow Y$ is a *closed immersion*, i.e. $X = \text{Spec}(B/\mathfrak{b})$ for an ideal \mathfrak{b} of B . Since f_* and $[Nf]_*$ are exact in this case (note that Nf is a **closed immersion* by 4.4), we only have to show $Nf_*\tilde{M} \cong [Nf]_*[N\tilde{M}]$ for all B/\mathfrak{b} -modules M of finite type or — equivalently — that $[Nf_*\tilde{M}](NY) \cong [[Nf]_*[N\tilde{M}]](NY)$. Now

$$[Nf_*\tilde{M}](NY) \stackrel{5.11}{\cong} [f_*\tilde{M}](Y) \otimes_B NB = \tilde{M}(X) \otimes_B NB = M \otimes_B NB$$

and (since $B \rightarrow C := B/\mathfrak{b}$ is a *finite ring homomorphism*)

$$\begin{aligned} [[Nf]_*[N\tilde{M}]](NY) &= [N\tilde{M}](NX) \stackrel{5.11}{\cong} \tilde{M}(X) \otimes_C NC \\ &= M \otimes_C NC \stackrel{4.16}{\cong} M \otimes_C (C \otimes_B NB) \cong M \otimes_B NB, \end{aligned}$$

so the theorem is true for closed immersions as well.

As a **third case**, we take an arbitrary *projective* morphism $f : X \rightarrow Y$. Since Y is affine, it admits an ample bundle, which implies (see [EGA II, 5.5.4(ii)]) that there is a $d \in \mathbb{N}_0$ for which f factorizes as $X \xrightarrow{i} \mathbb{P}_Y^d \xrightarrow{\pi} Y$, where i is a closed immersion and π is the structural morphism. Then for every coherent \mathcal{O}_X -modules \mathcal{F} and every $n \in \mathbb{Z}$, we have (because i_* and $[Ni]_*$ are exact)

$$\begin{aligned} NR^n f_* \mathcal{F} &= NR^n [\pi i]_* \mathcal{F} = NR^n \pi_* i_* \mathcal{F} \stackrel{1.\text{case}}{\cong} R^n [N\pi]_* N i_* \mathcal{F} \\ &\stackrel{2.\text{case}}{\cong} R^n [N\pi]_* [Ni]_* N \mathcal{F} = R^n [Nf]_* N \mathcal{F}, \end{aligned}$$

and the proof of this case is complete.

Finally we consider the **general case** of an arbitrary *proper* morphism $f : X \rightarrow Y$ and imitate Grothendieck's proof of the finiteness theorem for coherent modules [EGA III₁, 3.2.1]. Consider the full subcategory \mathcal{C} of Coh_X consisting of those coherent sheaves for which the theorem holds. We claim that \mathcal{C} has the following properties:

- (i) \mathcal{C} is *exact*, i.e. if $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ is a short exact sequence in Coh_X and if two of the three sheaves \mathcal{F}' , \mathcal{F} and \mathcal{F}'' belong to \mathcal{C} , then so does the third (compare [EGA III₁, 3.1.1]).
- (ii) If a coherent \mathcal{O}_X -module \mathcal{F} belongs to \mathcal{C} , then every direct factor of \mathcal{F} also belongs to \mathcal{C} .

Let $0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0$ be a short exact sequence as in (i). Applying the morphism of δ -functors (24), we get the following commutative diagram with exact rows

$$\begin{array}{ccccccccc} \cdots & \longrightarrow & NR^{n-1} f_* \mathcal{F}'' & \xrightarrow{\delta} & NR^n f_* \mathcal{F}' & \longrightarrow & NR^n f_* \mathcal{F} & \longrightarrow & NR^n f_* \mathcal{F}'' & \xrightarrow{\delta} & NR^{n+1} f_* \mathcal{F}' & \longrightarrow & \cdots \\ & & \downarrow \gamma_{n-1} & & \downarrow \alpha_n & & \downarrow \beta_n & & \downarrow \gamma_n & & \downarrow \alpha_{n+1} & & \\ \cdots & \longrightarrow & R^{n-1} [Nf]_* N \mathcal{F}'' & \xrightarrow{\delta} & R^n [Nf]_* N \mathcal{F}' & \longrightarrow & R^n [Nf]_* N \mathcal{F} & \longrightarrow & R^n [Nf]_* N \mathcal{F}'' & \xrightarrow{\delta} & R^{n+1} [Nf]_* N \mathcal{F}' & \longrightarrow & \cdots \end{array}$$

If two of \mathcal{F}' , \mathcal{F} and \mathcal{F}'' belong to \mathcal{C} , then for every n , two of α_n , β_n and γ_n are isomorphisms. The five lemma shows that then all α_n , β_n and γ_n are isomorphisms and hence \mathcal{F}' , \mathcal{F} and \mathcal{F}'' all belong to \mathcal{C} , which proves (i).

For (ii), let \mathcal{F} be a coherent \mathcal{O}_X -module in \mathcal{C} , and let \mathcal{F}_1 be a direct factor of \mathcal{F} . Putting $\mathcal{F}_2 := \mathcal{F} / \mathcal{F}_1$, we get a split short exact sequence

$$0 \longrightarrow \mathcal{F}_1 \longrightarrow \mathcal{F} = \mathcal{F}_1 \oplus \mathcal{F}_2 \longrightarrow \mathcal{F}_2 \longrightarrow 0$$

and hence for any n a morphism of split short exact sequences

$$\begin{array}{ccccccc}
0 & \longrightarrow & N R^n f_* \mathcal{F}_1 & \longrightarrow & N R^n f_* \mathcal{F} = N R^n f_* \mathcal{F}_1 \oplus N R^n f_* \mathcal{F}_2 & \longrightarrow & N R^n f_* \mathcal{F}_1 \longrightarrow 0 \\
& & \downarrow \alpha_n & & \downarrow \wr \alpha_n \oplus \beta_n & & \downarrow \beta_n \\
0 & \longrightarrow & R^n[Nf]_* N \mathcal{F}_1 & \longrightarrow & R^n[Nf]_* N \mathcal{F} = R^n[Nf]_* N \mathcal{F}_1 \oplus R^n[Nf]_* N \mathcal{F}_2 & \longrightarrow & R^n[Nf]_* N \mathcal{F}_1 \longrightarrow 0
\end{array}$$

with an isomorphism $\alpha_n \oplus \beta_n$ (because \mathcal{F} is in \mathcal{C}). It follows immediately that α_n and β_n must also be isomorphisms, i.e. \mathcal{F}_1 and \mathcal{F}_2 also belong to \mathcal{C} , which proves (ii).

In order to finish the proof of the theorem, we have to show that every coherent \mathcal{O}_X -module belongs to \mathcal{C} , and we want to do so by using *déviissage*: By [EGA III₁, 3.1.3], a full subcategory \mathcal{C} of Coh_X satisfying (i) and (ii) contains *all* coherent \mathcal{O}_X -modules if (and only if) for every irreducible closed subscheme Z of X , there is a sheaf with support Z in \mathcal{C} .

Let $Z \xrightarrow{i} X$ be a closed immersion with Z irreducible. Assume that we have found a coherent sheaf \mathcal{F}_Z of \mathcal{O}_Z -modules with support Z such that the theorem holds for \mathcal{F}_Z and the (obviously proper) morphism $f \circ i : Z \rightarrow Y$. Then $\mathcal{F} := i_* \mathcal{F}_Z$ is a coherent sheaf of \mathcal{O}_X -modules with support Z , and

$$N R^n f_* \mathcal{F} \cong N R^n [fi]_* \mathcal{F}_Z \cong R^n[N(fi)]_* N \mathcal{F}_Z \cong R^n[Nf]_* [(Ni)_* N \mathcal{F}_Z] \stackrel{2.\text{case}}{\cong} R^n[Nf]_* N \mathcal{F},$$

i.e. \mathcal{F} belongs to \mathcal{C} . Thus without loss of generality, we only have to consider the case $Z = X$ and therefore must exhibit a sheaf in \mathcal{C} with support X .

By Chow's lemma [EGA II, 5.6.2], there is a projective and surjective morphism $g : X' \rightarrow X$, with X' irreducible, such that the composition $f \circ g : X' \rightarrow Y$ is projective. Let $\mathcal{O}_{X'}(1)$ be a very ample bundle for g . Then by [EGA III₁, 2.2.1] and [EGA II, 3.4.7], there is an $m \in \mathbb{N}_0$ such that $\mathcal{F} := g_* \mathcal{O}_{X'}(m)$ has support X and such that

$$(25) \quad \forall n > 0 : R^n g_* \mathcal{O}_{X'}(m) = 0.$$

From (25), we learn two things. First, using the spectral sequence $R^p f_* R^q g_* \mathcal{O}_{X'}(m) \Rightarrow R^{p+q} [fg]_* \mathcal{O}_{X'}(m)$, we get

$$(26) \quad \forall n \in \mathbb{Z} : R^n f_* \mathcal{F} \cong R^n [fg]_* \mathcal{O}_{X'}(m).$$

Second, applying the third case to $\mathcal{O}_{X'}(m)$ and g , we get

$$\forall n > 0 : R^n [Ng]_* N \mathcal{O}_{X'}(m) \cong N R^n g_* \mathcal{O}_{X'}(m) \stackrel{(25)}{=} 0,$$

and then, using the spectral sequence $R^p [Nf]_* R^q [Ng]_* N \mathcal{O}_{X'}(m) \Rightarrow R^{p+q} [N(fg)]_* N \mathcal{O}_{X'}(m)$,

$$(27) \quad \forall n \in \mathbb{Z} : R^n [Nf]_* N \mathcal{F} \cong R^n [N(fg)]_* N \mathcal{O}_{X'}(m).$$

Combining these and applying the third case again, this time to $\mathcal{O}_{X'}(m)$ and fg , we get

$$N R^n f_* \mathcal{F} \stackrel{(26)}{\cong} N R^n [fg]_* \mathcal{O}_{X'}(m) \stackrel{3.\text{case}}{\cong} R^n [N(fg)]_* N \mathcal{O}_{X'}(m) \stackrel{(27)}{\cong} R^n [Nf]_* N \mathcal{F}$$

for all $n \in \mathbb{Z}$, i.e. \mathcal{F} belongs to \mathcal{C} , and the proof of the theorem is complete.

q.e.d.

7.6. Corollary. If k is a $*$ field and if X is a proper k -scheme, we have a canonical isomorphism

$$H^n(X, \mathcal{F}) \xrightarrow{\sim} H^n(NX, N\mathcal{F})$$

of finite dimensional k -vector spaces for every coherent \mathcal{O}_X -module \mathcal{F} and every $n \in \mathbb{N}_0$,

Proof. This follows immediately from 7.5, applied to $f : X \rightarrow \text{Spec}(k)$, and from 5.10:

$$\begin{aligned}
H^n(X, \mathcal{F}) &= H^n(X, \mathcal{F}) \otimes_k k = H^n(X, \mathcal{F}) \otimes_k Nk \\
&\stackrel{5.10}{=} N_* N R^n f_* \mathcal{F} \stackrel{7.5}{=} N_* R^n [Nf]_* N \mathcal{F} = H^n(NX, N\mathcal{F}).
\end{aligned}$$

q.e.d.

7.7. Corollary. For a * field k and a proper k -scheme X , the functor $N : \mathcal{Coh}_X \longrightarrow {}^*\mathcal{Coh}_{NX}$ is exact and fully faithful.

Proof. We already know that N is exact (and faithful) from 6.4(iv), even if X is not proper over k .

If $f : X \rightarrow \mathrm{Spec}(k)$ is proper, and if \mathcal{F} and \mathcal{G} are coherent \mathcal{O}_X -modules, we have

$$\begin{aligned} \mathrm{Hom}_{\mathcal{O}_{NX}}(N\mathcal{F}, N\mathcal{G}) &= \left[\mathcal{H}om_{\mathcal{O}_{NX}}(N\mathcal{F}, N\mathcal{G}) \right] (NX) \stackrel{6.5}{\cong} \left[N\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}) \right] (NX) \\ &= H^0(NX, N\mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})) \stackrel{7.6}{\cong} H^0(X, \mathcal{H}om_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G})) = \mathrm{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{G}), \end{aligned}$$

which proves fully faithfulness. **q.e.d.**

7.8. Corollary. For a * field k and a proper k -scheme X , the canonical group homomorphism $N : \mathrm{Pic}(X) \longrightarrow {}^*\mathrm{Pic}(NX)$ from 5.15 is *injective*.

Proof. This follows immediately from the fact that $N : \mathcal{Mod}_X^{\mathrm{fp}} \longrightarrow {}^*\mathcal{Mod}_{NX}^{\mathrm{fp}}$ is fully faithful by 7.7. **q.e.d.**

7.9. Example. Let k be a * field, and consider projective d -space over k for a $d \in \mathbb{N}_+$. Then the monomorphism $\mathrm{Pic}(\mathbb{P}_k^d) \hookrightarrow \mathrm{Pic}({}^*\mathbb{P}_k^d)$ from 7.8 is explicitly given by the following commutative diagram of abelian groups:

$$\begin{array}{ccccc} m & \in & \mathbb{Z} & \xrightarrow{\quad * \quad} & {}^*\mathbb{Z} & \ni & m \\ \downarrow & & \downarrow \wr & & \downarrow \wr & & \downarrow \\ \mathcal{O}_{\mathbb{P}_k^d}(m) & \in & \mathrm{Pic}(\mathbb{P}_k^d) & \xrightarrow{\quad N \quad} & {}^*\mathrm{Pic}({}^*\mathbb{P}_k^d) & \ni & \mathcal{O}_{{}^*\mathbb{P}_k^d}(m) \end{array}$$

7.10. Corollary. Let X be proper over a * field k , and let \mathcal{F} be a coherent \mathcal{O}_X -module. Then $\chi(\mathcal{F})$, the Euler-Poincaré characteristic of \mathcal{F} , equals $\chi(N\mathcal{F})$, the * Euler-Poincaré characteristic of $N\mathcal{F}$.

Proof. We have

$$\begin{aligned} \chi(N\mathcal{F}) &= \sum_{n=0}^{*\dim(NX)} (-1)^n \cdot {}^*\dim \left[H^n(NX, N\mathcal{F}) \right] \\ &\stackrel{6.4(ii)}{=} \sum_{n=0}^{\dim X} (-1)^n \cdot {}^*\dim \left[H^n(NX, N\mathcal{F}) \right] \stackrel{7.6}{=} \sum_{n=0}^{\dim X} (-1)^n \cdot \dim \left[H^n(X, \mathcal{F}) \right] = \chi(\mathcal{F}). \end{aligned}$$

q.e.d.

7.11. Corollary. Let X be a k -scheme of finite type, and let \mathcal{I} and \mathcal{J} be two sheaves of ideals in \mathcal{O}_X . Then $N[\mathcal{I} \cdot \mathcal{J}] = [N\mathcal{I}] \cdot [N\mathcal{J}]$ as * ideals of \mathcal{O}_{NX} .

Proof. Let Z be the closed subscheme of X given by $\mathcal{I} \cdot \mathcal{J}$, and let $i : Z \hookrightarrow X$ be the corresponding closed immersion. Then we have an exact sequence of coherent \mathcal{O}_X -modules

$$\mathcal{I} \otimes_{\mathcal{O}_X} \mathcal{J} \longrightarrow \mathcal{O}_X \longrightarrow i_*\mathcal{O}_Z \longrightarrow 0$$

and hence by 5.4, 5.14, 6.4(iv) and 7.5 an exact sequence

$$[N\mathcal{I}] \otimes_{\mathcal{O}_{NX}} [N\mathcal{J}] \xrightarrow{\varphi} \mathcal{O}_{NX} \longrightarrow [Ni]_*\mathcal{O}_{NZ} \longrightarrow 0$$


of * coherent \mathcal{O}_{NX} - * modules. By transfer, the image of φ is $[N\mathcal{I}] \cdot [N\mathcal{J}]$ and the ideal defining NZ , which in turn is $N[\mathcal{I} \cdot \mathcal{J}]$ by 6.6. **q.e.d.**

8. THE SHADOW MAP

Let $\langle K, |\cdot| : K \rightarrow \mathbb{R}^{\geq 0} \rangle$ be a non-trivially valued field with locally compact completion $\langle \hat{K}, |\cdot| \rangle$. Examples of such fields are \mathbb{Q} , \mathbb{R} and \mathbb{C} with their usual absolute value, \mathbb{Q}_p or \mathbb{Q}_p , equipped with the p -adic value $|\cdot|_p$ for a prime p or — more generally — local fields.

Assume that $\langle \hat{K}, |\cdot| \rangle$ is an element of our superstructure \hat{M} (which is no restriction, since we can always choose an appropriately large M).

Then $\langle {}^*K, |\cdot| \rangle$ and $\langle {}^*\hat{K}, |\cdot| \rangle$ are elements of ${}^*\hat{M}$, where ${}^*K \subseteq {}^*\hat{K}$ are fields, and $|\cdot| : {}^*K \rightarrow {}^*\mathbb{R}^{\geq 0}$ and $|\cdot| : {}^*\hat{K} \rightarrow {}^*\mathbb{R}^{\geq 0}$ are maps such that



commutes. By transfer we have

$$(M1) \quad \forall x \in {}^*\hat{K} : |x| = 0 \iff x = 0,$$

$$(M2) \quad \forall x, y \in {}^*\hat{K} : |x \cdot y| = |x| \cdot |y| \text{ and}$$

$$(M3) \quad \forall x, y \in {}^*\hat{K} : |x + y| \leq |x| + |y|.$$

Define the set of *finite* elements of *K by

$${}^*K^{\text{fin}} := \{x \in {}^*K \mid \exists C \in \mathbb{R}^{\geq 0} : |x| < C\}$$

and the set of *infinitesimal* elements of *K by

$${}^*K^{\text{inf}} := \{x \in {}^*K \mid \forall \varepsilon \in \mathbb{R}^{>0} : |x| < \varepsilon\}.$$

8.1. Proposition. ${}^*K^{\text{fin}} \subsetneq {}^*K$ is a valuation ring with maximal ideal ${}^*K^{\text{inf}}$ and residue field canonically isomorphic to \hat{K} . We call the projection ${}^*K^{\text{fin}} \twoheadrightarrow \hat{K}$ the *shadow map*, denote it by sh , and consequently get a commutative diagram of ring homomorphisms with exact row

(28)

$$\begin{array}{ccccccc} & & & & K & & \\ & & & & \downarrow & & \\ 0 & \longrightarrow & {}^*K^{\text{inf}} & \longrightarrow & {}^*K^{\text{fin}} & \xrightarrow{\text{sh}} & \hat{K} \longrightarrow 0 \\ & & & & \downarrow & & \\ & & & & {}^*K & & \end{array}$$

Proof. (M2) and (M3) immediately imply that ${}^*K^{\text{fin}}$ is a subring of *K . Since the value on K is non-trivial, the set of values is not bounded, so by transfer *K contains elements of infinite value, and ${}^*K^{\text{fin}}$ is a proper subring of *K .

If $x \in {}^*K$ is not finite, it in particular satisfies $|x| > 1$. Then $|\frac{1}{x}| < 1$ (by (M2)), i.e. $\frac{1}{x}$ is finite. This proves that ${}^*K^{\text{fin}}$ is indeed a valuation ring.

For a finite $x \in {}^*K^{\text{fin}} \setminus \{0\}$, $\frac{1}{x}$ is obviously infinite if and only if x is infinitesimal, which shows that ${}^*K^{\text{inf}}$ is the maximal ideal of ${}^*K^{\text{fin}}$.

Choose an infinite natural number h . We define a ring homomorphism $\alpha : \hat{K} \longrightarrow {}^*K^{\text{fin}}/{}^*K^{\text{inf}}$ by sending the class of a Cauchy sequence (x_n) in K to x_h . This is well-defined, because Cauchy sequences

are bounded (so that $x_h \in {}^*K^{\text{fin}}$) and because $\lim_{n \rightarrow \infty} x_n = 0$ implies $x_h \in {}^*K^{\text{inf}}$. Furthermore, α does not depend on h : If h' is another infinite natural number, and if (x_n) is a Cauchy sequence in K , then $x_h - x_{h'}$ is infinitesimal. Since \hat{K} is a field, α is automatically injective.

To prove that it is also surjective, we need the fact that \hat{K} is locally compact: This fact implies that there exists an $\varepsilon \in \mathbb{R}^{>0}$ and a compact subset A of \hat{K} such that

$$U_\varepsilon(0, \hat{K}) := \{x \in \hat{K} \mid |x| < \varepsilon\} \subseteq A.$$

Now let x be an arbitrary element of ${}^*K^{\text{fin}}$, let $C \in \mathbb{R}^{\geq 0}$ with $|x| < C$, let $\pi \in K$ with $|\pi| > 1$, and let $\mathbf{n} \in \mathbb{N}_+$ with $|\pi^{\mathbf{n}}| = |\pi|^{\mathbf{n}} > \frac{C}{\varepsilon}$. Because multiplication by $\pi^{\mathbf{n}}$ is a homeomorphism from \hat{K} to itself, $B := \pi^{\mathbf{n}} A$ is also compact, and we have

$$U_C(0, \hat{K}) \subseteq U_{\varepsilon|\pi^{\mathbf{n}}|}(0, \hat{K}) \subseteq B$$

and hence

$$x \in \{y \in {}^*K \mid |y| < C\} = {}^*U_C(0, \hat{K}) \subseteq {}^*B \subseteq {}^*K.$$

According to the nonstandard characterization of compactness, applied to B , any element of *B is infinitesimally close to an element of B , so there is an \hat{x} in \hat{K} with $x - \hat{x} \in {}^*K^{\text{inf}}$, i.e. $x = \alpha(\hat{x})$. **q.e.d.**

8.2. Corollary. Let X be a proper scheme over *K . Then the canonical map $X({}^*K^{\text{fin}}) \rightarrow X({}^*K)$ is bijective.

Proof. This follows immediately from 8.1 and the valuative criterion of properness [Har93, II.4.7]. **q.e.d.**

8.3. Corollary. Let X be a proper scheme over K . Then there is a canonical shadow map $\text{sh}_X : [{}^*X]({}^*K) \rightarrow X(\hat{K})$, induced by $\text{sh} : {}^*K^{\text{fin}} \rightarrow \hat{K}$, such that the following diagram commutes:

$$\begin{array}{ccc} & X(K) & \\ & \downarrow & \\ & X(\hat{K}) & \\ & \uparrow \text{sh}_X & \\ X({}^*K) & \xrightarrow[\sim]{4.13} & [{}^*X]({}^*K). \end{array}$$

Proof. Applying the functor $X(-)$ to (28), we get the following commutative diagram, in which α is bijective by 8.2, so that we can define sh_X as $(\text{sh} \circ \alpha^{-1} \circ \beta^{-1})$:

$$\begin{array}{ccc} & X(K) & \\ & \downarrow & \\ X({}^*K^{\text{fin}}) & \xrightarrow{\text{sh}} & X(\hat{K}) \\ \downarrow \wr \alpha & & \uparrow \text{sh}_X \\ X({}^*K) & \xrightarrow[\sim]{\beta} & [{}^*X]({}^*K). \end{array}$$

q.e.d.

8.4. Example. Let $X \subseteq \mathbb{P}_K^d$ be a projective variety over K , and let $x = (x_0 : \dots : x_d)$ be a *K -valued point of *X . Put $C := \max\{|x_0|, \dots, |x_d|\} \in {}^*\mathbb{R}^{>0}$. Then

$$\text{sh}_X(x) = \left(\text{sh} \left[\frac{x_0}{C} \right] : \dots : \text{sh} \left[\frac{x_d}{C} \right] \right) \in X(\hat{K}) \subseteq \mathbb{P}_K^d(\hat{K}).$$

9. RESOLUTION OF SINGULARITIES AND WEAK FACTORIZATION

For us, a *variety* over a field k is an integral, separated k -scheme of finite type. Similarly, if k is internal, a **variety* over k is a **integral*, **separated* **scheme* in ${}^*Sch_k^{\text{fp}}$.

9.1. Lemma. Let k be a **field* in ${}^*\mathcal{R}$, and let X be a k -variety. Then NX is a k -**variety*.

Proof. This follows immediately from 4.4 and 6.4(iii). **q.e.d.**

Let k be a field, and let X be a projective k -variety. Then for us, a *resolution (of singularities)* of X is a proper, birational k -morphism $X' \rightarrow X$, where X' is a projective, smooth k -variety.

9.2. Proposition. Let k be a **field* in ${}^*\mathcal{R}$ of *external* characteristic zero, let $n \in \mathbb{N}_+$, and let X be a **projective* k -**variety* which admits a **closed* embedding into ${}^*\mathbb{P}_k^n$ of finite **degree*. Then there exists a **resolution* $f : X' \rightarrow X$ of X .

Proof. By 6.22, there is a projective k -variety Y with $NY = X$, and by Hironaka's celebrated result on resolutions of singularities in characteristic zero, there exists a resolution $g : Y' \rightarrow Y$ of Y .

Then $X' := NY'$ is a **projective*, **smooth* k -**variety* by 4.4, 4.14 and 9.1, and $f := Ng : X' \rightarrow X$ is **proper* and **birational* by 4.4 and 6.28. **q.e.d.**

Using 9.2, we can now easily give a conceptual proof of the following classical result of Eklof (see [Ekl69]):

9.3. Corollary. For any pair (n, d) of natural numbers, there exists a bound $C \in \mathbb{N}_+$, such that for any field k of characteristic $p \geq C$ and any closed subvariety X of \mathbb{P}_k^n of degree d , there exists a resolution of singularities of X .

Proof. Assume the statement is false. Then for every $i \in \mathbb{N}_+$, we find a field k_i of characteristic $p_i \geq i$ and a closed subvariety X_i of $\mathbb{P}_{k_i}^n$ of degree d which does not admit a resolution.

We then take the full subcategory of *Rings* with objects $(k_i)_{i \in \mathbb{N}_+}$ as our base category \mathcal{B} , choose an infinite $j \in {}^*\mathbb{N}$ and get a **field* k_j of **characteristic* $p_j \geq j$ in ${}^*\mathcal{R}$ and a **closed* **subvariety* X_j of ${}^*\mathbb{P}_{k_j}^n$ of **degree* d which does not admit a **resolution*.

But since p_j is infinite, the external characteristic of k_j is zero, and 9.2 states that there can be no such X_j . Thus our assumption leads to a contradiction, and the corollary is proven. **q.e.d.**

9.4. Definition. Let k be a field, let U be an open subscheme of a projective k -variety X , and let $n \in \mathbb{N}_0$ be a natural number. We say that U has *complexity* n if $X \setminus U$, equipped with its reduced structure, has at most n irreducible components and if all those components have degree at most n .

9.5. Lemma. Let k be **field* in ${}^*\mathcal{R}$, let X be a projective k -variety, and let U' be a **open* subscheme of NX of *finite* **complexity*. Then there is an open subscheme U of X with $NU = U'$.

Proof. By definition of complexity, there is an $n \in \mathbb{N}_0$, such that $[NX] \setminus U' = Z'_1 \cup \dots \cup Z'_n$ with **integral* **closed* **subschemes* Z'_i of NX of **degree* at most n , and by 6.21, there exist integral closed subschemes Z_1, \dots, Z_n of X with $NZ_i = Z'_i$ for all i . Put $U := X \setminus \bigcup_{i=1}^n Z_i = \bigcap_{i=1}^n [X \setminus Z_i]$. Then

$$NU = N \left(\bigcap_{i=1}^n [X \setminus Z_i] \right) \stackrel{4.1(i)}{=} \bigcap_{i=1}^n N[X \setminus Z_i] \stackrel{4.8}{=} \bigcap_{i=1}^n ([NX] \setminus [NZ_i]) = \bigcap_{i=1}^n ([NX] \setminus Z'_i) = U'.$$

q.e.d.

9.6. Definition. Let $\Phi : X \dashrightarrow Y$ be a birational map between proper nonsingular varieties over a field k , and let $U \subseteq X$ be an open subscheme where Φ is an isomorphism. Then a *weak factorization of Φ with respect to U* is a factoring of Φ into a sequence of blow-ups and blow-downs with nonsingular irreducible centers disjoint from U . The *length* of a weak factorization is the number of blow-ups and blow-downs in the sequence.

9.7. Lemma. Let k be a \ast field in $\ast\mathcal{R}$, let $\Phi : X \rightarrow Y$ be a birational morphism between proper, smooth k -varieties, and let $U \subseteq X$ be an open subscheme where Φ is an isomorphism. If Φ admits a weak factorization with respect to U of length n , then $N\Phi : NX \rightarrow NY$ admits a \ast weak \ast factorization with respect to NU of \ast length n .

Proof. The statement makes sense, because NX and NY are \ast proper, \ast nonsingular k - \ast varieties by 4.4, 4.14 and 9.1, $N\Phi$ is \ast birational by 6.28, and $[N\Phi]|_{NU}$ is trivially an isomorphism.

Furthermore, it follows immediately from 4.14, 6.4(iii) and 6.9 that N maps any weak factorization of Φ with respect to U of length n to a \ast weak \ast factorization with respect to NU of \ast length n . **q.e.d.**

9.8. Proposition. Let k be a \ast algebraically closed \ast field in $\ast\mathcal{R}$ of *external* characteristic zero, let $n \in \mathbb{N}_+$, let X and Y be \ast projective, \ast nonsingular k -schemes which admit a \ast closed embedding into $\ast\mathbb{P}_k^n$ of finite \ast degree, let $\Phi : X \rightarrow Y$ be a \ast birational *morphism* of k - \ast schemes whose \ast graph has finite \ast degree, and let U be a \ast open \ast subscheme of X of finite \ast complexity where Φ is an isomorphism. Then Φ admits a \ast weak \ast factorization with respect to U of *finite* \ast length.

Proof. By 6.12, 6.22, 6.25, 6.28 and 9.5, there are projective, nonsingular k -varieties X' and Y' , a birational morphism $\Phi' : X' \rightarrow Y'$ and an open subscheme U' of X' , such that $NX' = X$, $NY' = Y$, $N\Phi' = \Phi$ and $NU' = U$. Since k is an algebraically closed field of characteristic zero and since $\Phi'|_{U'}$ is an isomorphism by 6.26, we know from [AKMW02, 0.1.1] that Φ' admits a weak factorization with respect to U' . The claim now follows immediately from 9.7. **q.e.d.**

9.9. Definition. Let k be a field. A *WF-datum over k* is a pair $\langle \Phi, U \rangle$, where $\Phi : X \rightarrow Y$ is a birational morphism between projective, nonsingular k -varieties and where U is an open subscheme of X where Φ is an isomorphism. A *weak factorization of $\langle \Phi, U \rangle$ (of length n)* is a weak factorization of Φ with respect to U of length n .

Let $N \in \mathbb{N}_0$ be a natural number. We say that the WF-datum $\langle \Phi, U \rangle$ has *complexity n* if X and Y are (isomorphic to) closed subschemes of \mathbb{P}_k^n of degree at most n , if the graph of Φ has degree at most n and if U has complexity n .

9.10. Corollary. For any $N \in \mathbb{N}_0$, there exists a bound $C \in \mathbb{N}_+$, such that for any algebraically closed field k of characteristic $p \geq C$, any WF-datum of complexity N has a weak factorization.

Proof. This follows from 9.8 in the same way as 9.3 follows from 9.2. **q.e.d.**

9.11. Corollary. For any $N \in \mathbb{N}_0$, there exists a bound $D \in \mathbb{N}_+$, such that for any algebraically closed field k of characteristic zero, any WF-datum of complexity N has a weak factorization of length at most D .

Proof. This, again, follows in the same way as 9.3 and 9.10, using the fact that the \ast weak \ast factorization whose existence is proven in 9.8 has *finite* \ast length. **q.e.d.**

REFERENCES

- [AKMW02] Dan Abramovich, Kalle Karu, Kenji Matsuki, and Jarosław Włodarczyk. Torification and Factorization of Birational Maps. *Journal of the American Mathematical Society*, 15(3):531–572, 2002.
- [BS05] Lars Brünjes and Christian Serpé. Enlargements of Categories. *Theory and Applications of Categories*, 14:357–398, 2005. <http://www.tac.mta.ca/tac/>.
- [EH00] David Eisenbud and Joe Harris. *The Geometry of Schemes*. Number 197 in Graduate Texts in Mathematics. Springer, New York, Berlin, Heidelberg, 2000.
- [Ekl69] Paul C. Eklof. Resolution of Singularities in Prime Characteristic for almost all Primes. *Transactions of the American Mathematical Society*, 146:429–438, 1969.
- [SGA 6] Alexander Grothendieck, P. Berthelot, and L. Illusie. *Théorie des intersections et théorème de Riemann-Roch*, volume 225 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1971. Séminaire de Géométrie Algébrique du Bois-Marie 1966–1967 (SGA 6), Dirigé par P. Berthelot, A. Grothendieck et L. Illusie. Avec la collaboration de D. Ferrand, J. P. Jouanolou, O. Jussila, S. Kleiman, M. Raynaud et J. P. Serre.
- [EGA I] Alexander Grothendieck. Éléments de géométrie algébrique: I. Le langage des schémas. *Publications mathématiques de l’I.H.É.S.*, 4, 1960.
- [EGA II] Alexander Grothendieck. Éléments de géométrie algébrique: II. Étude globale élémentaire de quelques classes de morphismes. *Publications mathématiques de l’I.H.É.S.*, 8, 1961.
- [EGA III₁] Alexander Grothendieck. Éléments de géométrie algébrique: III. Étude cohomologique des faisceaux cohérents, Première Partie. *Publications mathématiques de l’I.H.É.S.*, 11, 1961.
- [FGA] Alexander Grothendieck. *Fondements de la géométrie algébrique [Extraits du Séminaire Bourbaki, 1957–1962]*. Secrétariat mathématique, Paris, 1962.
- [EGA IV₁] Alexander Grothendieck. Éléments de géométrie algébrique: IV. Étude locale des schémas et des morphismes de schémas, Première Partie. *Publications mathématiques de l’I.H.É.S.*, 20, 1964.
- [EGA IV₂] Alexander Grothendieck. Éléments de géométrie algébrique: IV. Étude locale des schémas et des morphismes de schémas, Seconde Partie. *Publications mathématiques de l’I.H.É.S.*, 24, 1965.
- [EGA IV₃] Alexander Grothendieck. Éléments de géométrie algébrique: IV. Étude locale des schémas et des morphismes de schémas, Troisième Partie. *Publications mathématiques de l’I.H.É.S.*, 28, 1966.
- [EGA IV₄] Alexander Grothendieck. Éléments de géométrie algébrique: IV. Étude locale des schémas et des morphismes de schémas, Quatrième Partie. *Publications mathématiques de l’I.H.É.S.*, 32, 1967.
- [Har66] Robin Hartshorne. *Residues and Duality*, volume 20 of *Lecture Notes in Mathematics*. Springer, New York, Berlin, Heidelberg, 1966.
- [Har93] Robin Hartshorne. *Algebraic Geometry*, volume 52 of *Graduate Texts in Mathematics*. Springer, New York, Berlin, Heidelberg, 1993.
- [Kol96] János Kollár. *Rational Curves on Algebraic Surfaces*. Springer, Berlin, Heidelberg, New York, 1996.
- [MacL98] Saunders MacLane. *Categories for the Working Mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer, New York, Berlin, Heidelberg, 2nd edition, 1998.
- [Maci00] Angus Macintyre. Weil cohomology and model theory. In *Connections between model theory and algebraic and analytic geometry*, volume 6 of *Quaderni di Matematica*, pages 179–199. Dept. Math., Seconda Univ. Napoli, Caserta, 2000.
- [Mil80] J. S. Milne. *Étale Cohomology*. Princeton University Press, Princeton, New Jersey, 1980.
- [TT90] R. W. Thomason and Thomas Trobaugh. Higher Algebraic K-Theory of Schemes and of Derived Categories. In *The Grothendieck Festschrift III*, pages 247–435. Birkhäuser, Boston, 1990.
- [vdDS84] Lou van den Dries and K. Schmidt. Bounds in the theory of polynomial rings over fields. A nonstandard approach. *Inventiones mathematicae*, 76:77–91, 1984.

UNIVERSITÄT REGENSBURG, NWF I - MATHEMATIK, D-93040 REGENSBURG, GERMANY
 E-mail address: lars.bruejjes@mathematik.uni-regensburg.de

CHRISTIAN SERPÉ, WESTFÄLISCHE WILHELMS-UNIVERSITÄT MÜNSTER, MATHEMATISCHES INSTITUT, SONDERFORSCHUNGS-
 BEREICH 478 “GEOMETRISCHE STRUKTUREN IN DER MATHEMATIK”, HITTORFSTR. 27, D-48149 MÜNSTER, GERMANY
 E-mail address: serpe@math.uni-muenster.de